

characterized by an <sup>underlying</sup> ~~underlying~~ chronic inflammatory or ~~infectious~~ disease state (e.g., rheumatoid arthritis, osteomyelitis, tuberculosis, leprosy).

- 5 Heredofamilial amyloidoses may have associated neuropathic, renal, or cardiovascular deposits of the ATTR transthyretin type. Other heredofamilial amyloidoses include other syndromes and may have different amyloid components (e.g., familial Mediterranean fever which is characterized by AA fibrils). Other forms of amyloidosis include local forms, characterized by focal, often tumor-like deposits that occur in isolated organs. Other amyloidoses are associated with aging, and are commonly characterized by plaque formation in the heart or brain. Also common are amyloid deposits associated with long term hemodialysis. These and other forms of amyloid disease are summarized in Table 1. (Tan, S.Y. and Pepys, Histopathology 25:403-414, 1994; Harrison's Handbook of Internal Medicine, 13<sup>th</sup> Ed., Isselbacher, K.J., et al, eds, McGraw-Hill, San Francisco, 1995).
- 10

**Table 1**  
**Classification of Amyloid Diseases**

Amyloid Protein/Peptide	Protein Precursor	Protein Variants	Clinical
AA	Serum Amyloid A Protein (ApoSSA)		Reactive (secondary) Amyloidosis: Familial Mediterranean fever Familial amyloid nephropathy with urticaria and deafness (Muckle-Wells syndrome)
AA	Serum amyloid A protein (ApoSSA)		Reactive systemic amyloidosis associated with systemic inflammatory diseases
AL	Monoclonal immunoglobulin light chains (kappa, lambda)	Ak, A, (e.g., AkII)	Idiopathic (primary) Amyloidosis: myeloma or macroglobulinemia-associated; systemic amyloidosis associated with immunocyte

			dyscrasia; monoclonal gammopathy; occult dyscrasia; local nodular amyloidosis associated with chronic inflammatory diseases
AH	IgG (1( $\gamma$ 1))	A $\gamma$ 1	Heavy chain amyloidosis associated with several immunocyte dyscrasias
ATTR	Transthyretin (TTR)	At least 30 known point mutations	Familial amyloid polyneuropathy (e.g., Met 30, Portuguese)
ATTR	Transthyretin (TTR)	e.g., Met 111	Familial amyloid cardiomyopathy (Danish)
ATTR	Transthyretin (TTR)	Wild-type TTR or Ile 122	Systemic senile amyloidosis
AapoAI	ApoAI	Arg 26	Familial amyloid polyneuropathy
Agel	Gelsolin	Asn 187	Familial amyloidosis (Finnish)
Acys	Cystatin C	Gln 68	Hereditary cerebral hemorrhage with amyloidosis (Icelandic)
A $\beta$	Amyloid $\beta$ protein precursor (e.g. $\beta$ -APP <sub>695</sub> )	Various: Gln 618,	Alzheimer's disease Down's syndrome Hereditary cerebral hemorrhage amyloidosis (Dutch) Sporadic cerebral amyloid angiopathy Inclusion body myositis
AB <sub>2</sub> M	Beta <sub>2</sub> microglobulin		Associated with chronic hemodialysis
Acal	(Pro)calcitonin	(Pro)calcitonin	Medullary carcinoma of thyroid
AANF	Atrial natriuretic factor		Focal Senile Amyloidoses: Isolated atrial amyloid

amyloid		Brain	
A $\beta$	$\beta$ -amyloid precursor protein		<del>Brain</del>
SVEP <sup>a</sup>	-		Seminal vesicles
AB <sub>2</sub> M	Beta <sub>2</sub> microglobulin		Prostate
	Keratin		Primary localized cutaneous amyloid (macular, papular)
PrP	Prion precursor protein (33-35 kDa cellular form)	Scrapie protein 27-30 kDa	Sporadic Creutzfeldt-Jacob Disease Kuru (transmissible spongiform encephalopathies, prion diseases)
AIAPP	Islet amyloid polypeptide (IAPP)		Islets of Langerhans Diabetes type II, Insulinoma
Peptide hormones, fragments	e.g., precalcitonin		Exocrine amyloidosis, associated with APUDomas

<sup>a</sup>Seminal vesicle exocrine protein

Often, fibrils forming the bulk of an amyloid deposit are derived from one or more primary precursor proteins or peptides, and are usually associated with sulfated glycosaminoglycans. In addition, amyloid deposits may include minor proteins and peptides of various types, along with other components, such as proteoglycans, gangliosides and other sugars, as described in more detail in the sections that follow.

Currently, there are no specific, amyloid-directed treatments for any of the amyloid diseases. Where there is an underlying or associated disease state, therapy is directed towards decreasing the production of amyloidogenic protein by treating the underlying disease. This is exemplified by the treatment of tuberculosis with antibiotics, thereby reducing the mycobacterial load, resulting in a reduction of inflammation and in associated reduction of SSA protein. In the case of AL amyloid due to multiple myeloma, chemotherapy is administered to patients, causing a reduction in plasma cells and a lowering of myeloma immunoglobulin levels. As these levels decline, the AL amyloid may clear. Co-owned U.S. patent applications USSN 09/201,430, filed November 30, 1998 and USSN 09/322,289, filed May 28, 1999 reveal that amyloid plaque burden associated with Alzheimer's disease can be greatly reduced (and prevented) by administration of agents which produce or confer an immune response directed at  $\beta$ -

(A $\beta$ ) and

discovery

amyloid peptide (A $\beta$ ) and fragments thereof. It is the discovery of the present invention that induction of an immune response to various amyloid plaque components is effective in treating a broad range of amyloid diseases.

## 5 Summary of the Invention

The present invention is directed to pharmaceutical compositions and methods for treating a number of amyloid diseases. According to one aspect, the invention includes pharmaceutical compositions that include, as an active ingredient, an agent that is effective to induce an immune response against an amyloid component in a patient. Such  
 10 compositions will generally also include excipients and in preferred embodiments may include adjuvants. In further preferred embodiments, the adjuvants include, for example, aluminum hydroxide, aluminum phosphate, MPL<sup>TM</sup>, QS-21 (Stimulon<sup>TM</sup>) or incomplete Freund's adjuvant. According to a related embodiment, such pharmaceutical compositions may include a plurality of agents effective to induce an immune response against more  
 15 than one amyloid component in the patient.

In a related embodiment, the agent is effective to produce an immune response directed against a fibril peptide or protein amyloid component. Preferably, such a fibril peptide or protein is derived from a fibril precursor protein known to be associated with certain forms of amyloid diseases, as described herein. Such precursor proteins include,  
 20 but are not limited to, Serum Amyloid A protein (ApoSSA), immunoglobulin light chain, immunoglobulin heavy chain, ApoAI, transthyretin, lysozyme, fibrogen  $\alpha$  chain, gelsolin, cystatin C, Amyloid  $\beta$  protein precursor ( $\beta$ -APP), Beta<sub>2</sub> microglobulin, prion precursor protein (PrP), atrial natriuretic factor, keratin, islet amyloid polypeptide, a peptide hormone, and synuclein. Such precursors also include mutant proteins, protein fragments  
 25 and proteolytic peptides of such precursors. In a preferred embodiment, the agent is effective to induce an immune response directed against a neoepitope formed by a fibril protein or peptide, with respect to a fibril precursor protein. That is, as described in more detail herein, many fibril-forming peptides or proteins are fragments of such precursor proteins, such as those listed above. When such fragments are formed, such as by  
 30 proteolytic cleavage, epitopes may be revealed that are not present on the precursor and are therefore not immunologically available to the immune system when the fragment is a part



protein.

may be

of the precursor ~~prot-~~ Agents directed to such epitopes ~~may be~~ preferred therapeutic agents, since they may be less likely to induce an autoimmune response in the patient.

According to a related embodiment, pharmaceutical compositions of the invention include agents directed to amyloid components, such as those selected from the group including, but not limited to the following fibril peptides or proteins: AA, AL, ATTR, 5 AApoA1, Alys, Agel, Acys, A $\beta$ , AB<sub>2</sub>M, AScr, Acal, AIAPP and synuclein-NAC fragment. The full names and compositions of these peptides are described herein. Such peptides can be made according to methods well known in the art, as described herein.

According to a further related embodiment, agents included in such pharmaceutical 10 compositions also include certain to sulfated proteoglycans. In a related embodiment, the proteoglycan is a heparin sulfate glycosaminoglycan, preferably perlecan, dermatan sulfate, chondroitin-4-sulfate, or pentosan polysulfate.

According to another related aspect, the invention includes a method of preventing or treating a disorder characterized by amyloid deposition in a mammalian subject. In 15 accordance with this aspect of the invention, the subject is given a dosage of an agent effective to produce an immune response against an amyloid component characteristic of the amyloid disorder from which the subject suffers. Essentially, the methods include administering pharmaceutical compositions containing immunogenic amyloid components specific to the disorder, such as those described above. Such methods are further 20 characterized by their effectiveness in inducing immunogenic responses in the subject. According to a preferred embodiment, the method is effective to produce an immunological response that is characterized by a serum titer of at least 1:1000 with respect to the amyloid component against which the immunogenic agent is directed. In yet a further preferred embodiment, the serum titer is at least 1:5000 with respect to the fibril 25 component. According to a related embodiment, the immune response is characterized by a serum amount of immunoreactivity corresponding to greater than about four times higher than a serum level of immunoreactivity measured in a pre-treatment control serum sample. This latter characterization is particularly appropriate when serum immunoreactivity is measured by ELISA techniques, but can apply to any relative or absolute measurement of 30 serum immunoreactivity. According to a preferred embodiment, the immunoreactivity is measured at a serum dilution of about 1:100.

According to a still further related aspect, the invention includes a method of determining the prognosis of a patient undergoing treatment for an amyloid disorder.

amyloid

09/724570

amount

Here, patient serum ~~a~~<sup>amount</sup> of immunoreactivity against an ~~an~~<sup>amyloid</sup> component characteristic of the selected disorder is measured, and a patient serum amount of immunoreactivity of at least four times a baseline control level of serum immunoreactivity is indicative of a prognosis of improved status with respect to the particular amyloid disorder. According to preferred embodiments, the amount of immunoreactivity against the selected amyloid component present in the patient serum is characterized by a serum titer of at least about 1:1000, or at least 1:5000, with respect to the amyloid component.

According to a still related aspect, the invention also includes so-called "passive immunization" methods and pharmaceutical compositions for preventing or treating amyloid diseases. According to this aspect of the invention, patients are given an effective dosage of an antibody that specifically binds to a selected amyloid component, preferably a fibril component present in amyloid deposits characteristic of the disease to be treated. In general, such antibodies are selected for their abilities to specifically bind the various proteins, peptides, and components described with respect to the pharmaceutical compositions and methods described in the preceding paragraphs of this section. According to a related embodiment, such methods and compositions may include combinations of antibodies that bind at least two amyloid fibril components. In general, pharmaceutical compositions are administered to provide a serum amount of immunoreactivity against the target amyloid component that is at least about four times higher than a serum level of immunoreactivity against the component measured in a control serum sample. The antibodies may also be administered with a carrier, as described herein. In general, in accordance with this aspect of the invention, such antibodies, will be administered (or formulated for administration) peritoneally, orally, intranasally, subcutaneously, intramuscularly, topically or intravenously, but can be administered or formulated for administration by any pharmaceutically effective route (i.e., effective to produce the indicated therapeutic levels, as set forth above and herein).

According to a related embodiment, therapeutic antibodies may be administered by administering a polynucleotide encoding at least one antibody chain to the patient. According to this aspect of the invention, the polynucleotide is expressed in the patient to produce the antibody chain in a pharmaceutically effective amount in the patient. Such a polynucleotide may encode heavy and light chains of the antibody, thereby producing the heavy and light chains in the patient.

*preferred*

According to ~~referred~~ embodiments, the immunization regimens described above may include administration of agents, including antibodies, in multiple dosages, such as over a 6 month period, such as an initial immunization followed by booster injections at time intervals, such as 6 week intervals, according to methods known in the art, or  
 5 according to patient need, as assessed by immunological response. Alternatively, or in addition, such regimens may include the use of "sustained release" formulations, such as are known in the art.

These and other objects and features of the invention will become more fully apparent when the following detailed description of the invention is read in conjunction  
 10 with the accompanying drawings.

### Brief Description of the Figures

FIG. 1: Antibody titer in transgenic mice after injection with A $\beta$ 1-42.

FIG. 2: Amyloid burden in the hippocampus. The percentage of the area  
 15 of the hippocampal region occupied by amyloid plaques, defined by reactivity with the A $\beta$ -specific monoclonal antibody 3D6, was determined by computer-assisted quantitative image analysis of immunoreacted brain sections. The values for individual mice are shown sorted by treatment group. The horizontal line for each grouping indicates the median value of the distribution.

FIG. 3: Neuritic dystrophy in the hippocampus. The percentage of the area  
 20 of the hippocampal region occupied by dystrophic neurites, defined by their reactivity with the human APP-specific monoclonal 8E5, was determined by quantitative computer-assisted image analysis of immunoreacted brain sections. The values for individual mice are shown for the AN1792-treated group and the PBS-treated control group. The  
 25 horizontal line for each grouping indicates the median value of the distribution.

FIG. 4: Astrocytosis in the retrosplenial cortex. The percentage of the area  
 of the cortical region occupied by glial fibrillary acidic protein (GFAP)-positive astrocytes was determined by quantitative computer-assisted image analysis of immunoreacted brain sections. The values for individual mice are shown sorted by treatment group and median  
 30 group values are indicated by horizontal lines.

Geometric

42 following

FIG. 5. ~~Geometric~~ mean antibody titers to A $\beta$ 42 ~~following~~ immunization with a range of eight doses of A $\beta$ 42 ("AN1792") containing 0.14, 0.4, 1.2, 3.7, 11, 33, 100, or 300  $\mu$ g.

FIG. 6: Kinetics of antibody response to AN1792 immunization. Titers are expressed as geometric means of values for the 6 animals in each group.

FIG. 7: Quantitative image analysis of the cortical amyloid burden in PBS- and AN1792-treated mice.

FIG. 8: Quantitative image analysis of the neuritic plaque burden in PBS- and AN1792-treated mice.

FIG. 9: Quantitative image analysis of the percent of the retrosplenial cortex occupied by astrocytosis in PBS- and AN1792-treated mice.

FIG. 10: Lymphocyte Proliferation Assay on spleen cells from AN1792-treated (upper panel) or PBS-treated (lower panel) mice.

FIG. 11: Total A $\beta$  levels in the cerebral cortex. A scatterplot of individual A $\beta$  profiles in mice immunized with A $\beta$  or APP derivatives combined with Freund's adjuvant.

FIG. 12: Amyloid burden in the cortex, determined by quantitative image analysis of immunoreacted brain sections for mice immunized with the A $\beta$  peptide conjugates A $\beta$ 1-5, A $\beta$ 1-12, and A $\beta$ 13-28; the full length A $\beta$  aggregates A $\beta$ 42 ("AN1792") and A $\beta$ 1-40 ("AN1528") and the PBS-treated control group.

FIG. 13: Geometric mean titers of A $\beta$ -specific antibody for groups of mice immunized with A $\beta$  or APP derivatives combined with Freund's adjuvant.

FIG. 14: Geometric mean titers of A $\beta$ -specific antibody for groups of guinea pigs immunized with AN1792, or a palmitoylated derivative thereof, combined with various adjuvants.

FIG. 15 (A-E): A $\beta$  levels in the cortex of 12-month old PDAPP mice treated with AN1792 or AN1528 with different adjuvants.

FIG. 16: Mean titer of mice treated with polyclonal antibody to A $\beta$ .

FIG. 17: Mean titer of mice treated with monoclonal antibody 10D5 to A $\beta$ .

FIG. 18: Mean titer of mice treated with monoclonal antibody 2F12 to A $\beta$ .

## Detailed Description of the Invention

a chemical

biological

09/724570

An "agent" is ~~chemical~~ <sup>a chemical</sup> molecule of synthetic or ~~biological~~ <sup>biological</sup> origin. In the context of the present invention, an agent is generally a molecule that can be used in a pharmaceutical composition.

5 An "anti-amyloid agent" is an agent which is capable of producing an immune response against an amyloid plaque component in a vertebrate subject, when administered by active or passive immunization techniques.

10 The terms "polynucleotide" and "nucleic acid," as used interchangeably herein refer to a polymeric molecule having a backbone that supports bases capable of hydrogen bonding to typical polynucleotides, where the polymer backbone presents the bases in a manner to permit such hydrogen bonding in a sequence specific fashion between the polymeric molecule and a typical polynucleotide (e.g., single-stranded DNA). Such bases are typically inosine, adenosine, guanosine, cytosine, uracil and thymidine. Polymeric molecules include double and single stranded RNA and DNA, and backbone modifications thereof, for example, methylphosphonate linkages.

15 The term "polypeptide" as used herein refers to a compound made up of a single chain of amino acid residues linked by peptide bonds. The term "protein" may be synonymous with the term "polypeptide" or may refer to a complex of two or more polypeptides.

20 The term "peptide" also refers to a compound composed of amino acid residues linked by peptide bonds. Generally peptides are composed of 100 or fewer amino acids, while polypeptides or proteins have more than 100 amino acids. As used herein, the term "protein fragment" may also be read to mean a peptide.

25 A "fibril peptide" or "fibril protein" refers to a monomeric or aggregated form of a protein or peptide that forms fibrils present in amyloid plaques. Examples of such peptides and proteins are provided herein.

30 A "pharmaceutical composition" refers to a chemical or biological composition suitable for administration to a mammalian individual. Such compositions may be specifically formulated for administration via one or more of a number of routes, including but not limited to, oral, parenteral, intravenous, intraarterial, subcutaneous, intranasal, sublingual, intraspinal, intracerebroventricular, and the like.

A "pharmaceutical excipient" or a "pharmaceutically acceptable excipient" is a carrier, usually a liquid, in which an active therapeutic agent is formulated. The excipient generally does not provide any pharmacological activity to the formulation, though it may

and/or

characteristics,

provide chemical ~~and~~ biological stability, release ~~characteristics~~, and the like. Exemplary formulations can be found, for example, in Remington's Pharmaceutical Sciences, 19<sup>th</sup> Ed., Grennaro, A., Ed., 1995.

- 5 A "glycoprotein" is protein to which at least one carbohydrate chain (oligopolysaccharide) is covalently attached.

A "proteoglycan" is a glycoprotein where at least one of the carbohydrate chains is a glycosaminoglycan, which is a long linear polymer of repeating disaccharides in which one member of the pair usually is a sugar acid (uronic acid) and the other is an amino sugar.

- 10 The term "immunological" or "immune" or "immunogenic" response refers to the development of a humoral (antibody mediated) and/or a cellular (mediated by antigen-specific T cells or their secretion products) response directed against an antigen in a vertebrate individual. Such a response can be an active response induced by administration of immunogen or a passive response induced by administration of antibody
- 15 or primed T-cells. A cellular immune response is elicited by the presentation of polypeptide epitopes in association with Class I or Class II MHC molecules to activate antigen-specific CD4<sup>+</sup> T helper cells and/or CD8<sup>+</sup> cytotoxic T cells. The response may also involve activation of monocytes, macrophages, NK cells, basophils, dendritic cells, astrocytes, microglia cells, eosinophils or other components of innate immunity. The
- 20 presence of a cell-mediated immunological response can be determined by standard proliferation assays (CD4<sup>+</sup> T cells) or CTL (cytotoxic T lymphocyte) assays known in the art. The relative contributions of humoral and cellular responses to the protective or therapeutic effect of an immunogen can be distinguished by separately isolating immunoglobulin (IgG) and T-cell fractions from an immunized syngeneic animal and
- 25 measuring protective or therapeutic effect in a second subject.

*immunogenic*

An "~~immune~~<sup>immunogenic</sup> agent" or "immunogen" or "antigen" is a molecule that is capable of inducing an immunological response against itself upon administration to a patient, either in conjunction with, or in the absence of, an adjuvant. Such molecules include, for example, amyloid fibril peptides or fragments thereof conjugated to a carrier protein, such keyhole limpet hemocyanin, Cd3 or tetanus toxin.

An "epitope" or "antigenic determinate" is the part of an antigen that binds to the antigen-binding region of an antibody.

The term "A $\beta$ ," "A $\beta$  peptide" and "Amyloid  $\beta$ " peptide are synonymous, and refer to one or more peptide compositions of about 38-43 amino acids derived from Beta Amyloid Precursor Protein ( $\beta$ -APP), as described herein. "A $\beta$ xx" refers to amyloid  $\beta$  peptide 1-xx, where xx is a number indicating the number of amino acids in the peptide; e.g., A $\beta$ 42 is the same as A $\beta$ 1-42, which is also referred to herein as "AN1792," and A $\beta$ 40 is the same as A $\beta$ 1-40, which is also referred to herein as "AN1578."

Disaggregated or monomeric A $\beta$  means soluble, monomeric peptide units of A $\beta$ .  
 One method to prepare monomeric A $\beta$  is to dissolve lyophilized peptide in neat DMSO with sonication. The resulting solution is centrifuged to remove any insoluble particulates. Aggregated A $\beta$  is a mixture of oligomers in which the monomeric units are held together by noncovalent bonds.

The term "naked polynucleotide" refers to a polynucleotide not complexed with colloidal materials. Naked polynucleotides are sometimes cloned in a plasmid vector.

The term "patient" includes human and other mammalian subjects that receive either prophylactic or therapeutic treatment.

The terms "significantly different than," "statistically significant," "significantly higher (or lower) than," and similar phrases refer to comparisons between data or other measurements, wherein the differences between two compared individuals or groups are evidently or reasonably different to the trained observer, or statistically significant (if the phrase includes the term "statistically" or if there is some indication of statistical test, such as a p-value, or if the data, when analyzed, produce a statistical difference by standard statistical tests known in the art).

Compositions <sup>or</sup> methods "comprising" one or more recited elements may include other elements not specifically recited. For example, a composition that comprises a fibril component peptide encompasses both the isolated peptide and the peptide as a component of a larger polypeptide sequence. By way of further example, a composition that  
 5 comprises elements A and B also encompasses a composition consisting of A, B and C.

## B. Amyloid Diseases

### 1. Overview and Pathogenesis

Amyloid diseases or amyloidoses include a number of disease states having a wide  
 10 variety of outward symptoms. These disorders have in common the presence of abnormal extracellular deposits of protein fibrils, known as "amyloid deposits" or "amyloid plaques" that are usually about 10-100  $\mu$ m in diameter and are localized to specific organs or tissue regions. Such plaques are composed primarily of a naturally occurring soluble protein or peptide. These insoluble deposits are composed of generally lateral aggregates of fibrils  
 15 that are approximately 10-15 nm in diameter. Amyloid fibrils produce a characteristic apple green birefringence in polarized light, when stained with Congo Red dye. The disorders are classified on the basis of the major fibril components forming the plaque deposits, as discussed below.

The peptides or proteins forming the plaque deposits are often produced from a  
 20 larger precursor protein. More specifically, the pathogenesis of amyloid fibril deposits generally involves proteolytic cleavage of an "abnormal" precursor protein into fragments. These fragments generally aggregate into anti-parallel  $\beta$  pleated sheets; however, certain undegraded forms of precursor protein have been reported to aggregate and form fibrils in familial amyloid polyneuropathy (variant transthyretin fibrils) and dialysis-related  
 25 amyloidosis ( $\beta_2$  microglobulin fibrils) (Tan, *et al.*, 1994, *supra*).

### 2. Clinical Syndromes

This section provides descriptions of major types of amyloidoses, including their characteristic plaque fibril compositions. It is a general discovery of the present invention that amyloid diseases can be treated by administering agents that serve to stimulate an  
 30 immune response against a component or components of the various disease-specific amyloid deposits. As discussed in more detail in Section C below, such components are



constituents of

the

preferably ~~constitution~~ of the fibrils that form the plaques. ~~The~~ sections below serve to exemplify major forms of amyloidosis and are not intended to limit the invention.

a. AA (reactive) Amyloidosis

Generally, AA amyloidosis is a manifestation of a number of diseases that provoke a sustained acute phase response. Such diseases include chronic inflammatory disorders, chronic local or systemic microbial infections, and malignant neoplasms.

AA fibrils are generally composed of 8000 dalton fragments (AA peptide or protein) formed by proteolytic cleavage of serum amyloid A protein (apoSSA), a circulating apolipoprotein which is present in HDL particles and which is synthesized in hepatocytes in response to such cytokines as IL-1, IL-6 and TNF. Deposition can be widespread in the body, with a preference for parenchymal organs. The spleen is usually a deposition site, and the kidneys may also be affected. Deposition is also common in the heart and gastrointestinal tract.

AA amyloid diseases include, but are not limited to inflammatory diseases, such as rheumatoid arthritis, juvenile chronic arthritis, ankylosing spondylitis, psoriasis, psoriatic arthropathy, Reiter's syndrome, Adult Still's disease, Behçet's syndrome, and Crohn's disease. AA deposits are also produced as a result of chronic microbial infections, such as leprosy, tuberculosis, bronchiectasis, decubitus ulcers, chronic pyelonephritis, osteomyelitis, and Whipple's disease. Certain malignant neoplasms can also result in AA fibril amyloid deposits. These include such conditions as Hodgkin's lymphoma, renal carcinoma, carcinomas of gut, lung and urogenital tract, basal cell carcinoma, and hairy cell leukemia.

b. AL Amyloidoses

AL amyloid deposition is generally associated with almost any dyscrasia of the B lymphocyte lineage, ranging from malignancy of plasma cells (multiple myeloma) to benign monoclonal gammopathy. At times, the presence of amyloid deposits may be a primary indicator of the underlying dyscrasia.

Fibrils of AL amyloid deposits are composed of monoclonal immunoglobulin light chains or fragments thereof. More specifically, the fragments are derived from the N-terminal region of the light chain (kappa or lambda) and contain all or part of the variable ( $V_L$ ) domain thereof. Deposits generally occur in the mesenchymal tissues, causing peripheral and autonomic neuropathy, carpal tunnel syndrome, macroglossia, restrictive cardiomyopathy, arthropathy of large joints, immune dyscrasias, myelomas, as well as

However, it should be noted that almost any ~~tissue~~ <sup>tissue</sup>, particularly visceral organs such as the heart, may be involved.

c. Hereditary Systemic Amyloidoses

- There are many forms of hereditary systemic amyloidoses. Although they are relatively rare conditions, adult onset of symptoms and their inheritance patterns (usually autosomal dominant) lead to persistence of such disorders in the general population. Generally, the syndromes are attributable to point mutations in the precursor protein leading to production of variant amyloidogenic peptides or proteins. Table 2 summarizes the fibril composition of exemplary forms of these disorders.

10

Table 2  
Hereditary Amyloidoses<sup>a</sup>

Fibril Peptide/Protein	Genetic variant	Clinical Syndrome
Transthyretin and fragments (ATTR)	Met30, many others	Familial amyloid polyneuropathy (FAP), (mainly peripheral nerves)
Transthyretin and fragments (ATTR)	Thr45, Ala60, Ser84, Met111, Ile122	Cardiac involvement predominant without neuropathy
N-terminal fragment of Apolipoprotein A1 (apoA1)	Arg 26	Familial amyloid polyneuropathy (FAP), (mainly peripheral nerves)
N-terminal fragment of Apolipoprotein A1 (AapoA1)	Arg26, Arg50, Arg 60, others	Ostertag-type, non-neuropathic (predominantly visceral involvement)
Lysozyme (Alys)	Thr56, His67	Ostertag-type, non-neuropathic (predominantly visceral involvement)
Fibrogen $\alpha$ chain fragment	Leu554, Val 526	Ostertag-type, non-neuropathic (predominantly visceral involvement)
Gelsolin fragment (Agel)	Asn187, Tyr187	Cranial neuropathy with lattice corneal dystrophy
Cystatin C fragment	Glu68	Hereditary cerebral hemorrhage (cerebral amyloid angiopathy) – Icelandic type
$\beta$ -amyloid protein (A $\beta$ ) derived from Amyloid Precursor Protein (APP)	Gln693	Hereditary cerebral hemorrhage (cerebral amyloid angiopathy) – Dutch type
$\beta$ -amyloid protein (A $\beta$ ) derived from Amyloid Precursor Protein (APP)	Ile717, Phe717, Gly717	Familial Alzheimer's Disease
$\beta$ -amyloid protein (A $\beta$ ) derived from Amyloid	Asn670, Leu671	Familial Dementia – probable Alzheimer's Disease

APP

Precursor Protein (A <sub>2</sub> )		
Prion Protein (PrP) derived from PrP precursor protein 51-91 insert	Leu102, Val167, Asn178, Lys200	Familial Creutzfeldt-Jakob disease; Gerstmann-Sträussler-Scheinker syndrome (hereditary spongiform encephalopathies, prion diseases)
AA derived from Serum amyloid A protein (ApoSSA)		Familial Mediterranean fever, predominant renal involvement (autosomal recessive)
AA derived from Serum amyloid A protein (ApoSSA)		Muckle-Well's syndrome, nephropathy, deafness, urticaria, limb pain
Unknown		Cardiomyopathy with persistent atrial standstill
Unknown		Cutaneous deposits (bullous, papular, pustuloderma)

<sup>a</sup>Data derived from Tan & Pepys, 1994, *supra*.

The data provided in Table 2 are exemplary and are not intended to limit the scope of the invention. For example, more than 40 separate point mutations in the transthyretin gene have been described, all of which give rise to clinically similar forms of familial amyloid polyneuropathy.

Transthyretin (TTR) is a 14 kilodalton protein that is also sometimes referred to as prealbumin. It is produced by the liver and choroid plexus, and it functions in transporting thyroid hormones and vitamin A. At least 50 variant forms of the protein, each characterized by a single amino acid change, are responsible for various forms of familial amyloid polyneuropathy. For example, substitution of proline for leucine at position 55 results in a particularly progressive form of neuropathy; substitution of methionine for leucine at position 111 resulted in a severe cardiopathy in Danish patients. Amyloid deposits isolated from heart tissue of patients with systemic amyloidosis have revealed that the deposits are composed of a heterogeneous mixture of TTR and fragments thereof, collectively referred to as ATTR, the full length sequences of which have been characterized. ATTR fibril components can be extracted from such plaques and their structure and sequence determined according to the methods known in the art (e.g., Gustavsson, A., *et al.*, Laboratory Invest. 73: 703-708, 1995; Kametani, F., *et al.*, Biochem. Biophys. Res. Commun. 125: 622-628, 1984; Pras, M., *et al.*, PNAS 80: 539-42, 1983).

Persons <sup>having</sup> ~~having~~ point mutations in the molecule apolipoprotein AI (e.g., Gly→Arg26; Trp → Arg50; Leu → Arg60) exhibit a form of amyloidosis ("Östertag type") characterized by deposits of the protein apolipoprotein AI or fragments thereof (AApoAI). These patients have low levels of high density lipoprotein (HDL) and present  
 5 with a peripheral neuropathy or renal failure.

A mutation in the alpha chain of the enzyme lysozyme (e.g., Ile→Thr56 or Asp→His57) is the basis of another form of Östertag-type non-neuropathic hereditary amyloid reported in English families. Here, fibrils of the mutant lysozyme protein (Alys) are deposited, and patients generally exhibit impaired renal function. This protein, unlike  
 10 most of the fibril-forming proteins described herein, is usually present in whole (unfragmented) form (Benson, M.D., *et al.* CIBA Fdn. Symp. 199: 104-131, 1996).

β-amyloid peptide (Aβ) is a 39-43 amino acid peptide derived by proteolysis from a large protein known as beta amyloid precursor protein (βAPP). Mutations in βAPP result in familial forms of Alzheimer's disease, Down's syndrome and/or senile dementia,  
 15 characterized by cerebral deposition of plaques composed of Aβ fibrils and other components, which are described in further detail below. Known mutations in APP associated with Alzheimer's disease occur proximate to the cleavage sites of β or γ secretase, or within Aβ. For example, position 717 is proximate to the site of γ-secretase cleavage of APP in its processing to Aβ, and positions 670/671 are proximate to the site of  
 20 β-secretase cleavage. Mutations at any of these residues may result in Alzheimer's disease, presumably by causing an increase the amount of the 42/43 amino acid form of Aβ generated from APP. The structure and sequence of Aβ peptides of various lengths are well known in the art. Such peptides can be made according to methods known in the art (e.g., Glenner and Wong, Biochem Biophys. Res. Comm. 129: 885-890, 1984; Glenner  
 25 and Wong, Biochem Biophys. Res. Comm. 122: 1131-1135, 1984). In addition, various forms of the peptides are commercially available.

Synuclein is a synapse-associated protein that resembles an alipoprotein and is abundant in neuronal cytosol and presynaptic terminals. A peptide fragment derived from α-synuclein, termed NAC, is also a component of amyloid plaques of Alzheimer's disease.  
 30 (Clayton, *et al.*, 1998). This component also serves as a target for immunologically-based treatments of the present invention, as detailed below.

calcium

fragments

Gelsolin is a ~~calcium~~ binding protein that binds to and ~~fragments~~ actin filaments.

Mutations at position 187 (e.g., Asp→Asn; Asp→Tyr) of the protein result in a form of hereditary systemic amyloidosis, usually found in patients from Finland, as well as persons of Dutch or Japanese origin. In afflicted individuals, fibrils formed from gelsolin

- 5 fragments (Agel), usually consist of amino acids 173-243 (68 kDa carboxyterminal fragment) and are deposited in blood vessels and basement membranes, resulting in corneal dystrophy and cranial neuropathy which progresses to peripheral neuropathy, dystrophic skin changes and deposition in other organs. (Kangas, H., *et al.* Human Mol. Genet. 5(9): 1237-1243, 1996).

- 10 Other mutated proteins, such as mutant alpha chain of fibrinogen (AfibA) and mutant cystatin C (Acys) also form fibrils and produce characteristic hereditary disorders. AfibA fibrils form deposits characteristic of a nonneuropathic hereditary amyloid with renal disease; Acys deposits are characteristic of a hereditary cerebral amyloid angiopathy reported in Iceland. (Isselbacher, *et al.*, Harrison's Principles of
- 15 Internal Medicine, McGraw-Hill, San Francisco, 1995; Benson, *et al.*, *supra.*). In at least some cases, patients with cerebral amyloid angiopathy (CAA) have been shown to have amyloid fibrils containing a non-mutant form of cystatin C in conjunction with beta protein. (Nagai, A., *et al.* Molec. Chem. Neuropathol. 33: 63-78, 1998).

- Certain forms of prion disease are now considered to be heritable, accounting for
- 20 up to 15% of cases, which were previously thought to be predominantly infectious in nature. (Baldwin, *et al.*, in *Research Advances in Alzheimer's Disease and Related Disorders*, John Wiley and Sons, New York, 1995). In such prion disorders, patients develop plaques composed of abnormal isoforms of the normal prion protein (PrP<sup>C</sup>). A predominant mutant isoform, PrP<sup>Sc</sup>, also referred to as AScr, differs from the normal
- 25 cellular protein in its resistance to protease degradation, insolubility after detergent extraction, deposition in secondary lysosomes, post-translational synthesis, and high  $\beta$ -pleated sheet content. Genetic linkage has been established for at least five mutations resulting in Creutzfeldt-Jacob disease (CJD), Gerstmann-Sträussler-Scheinker syndrome (GSS), and fatal familial insomnia (FFI). (Baldwin) Methods for extracting fibril
- 30 peptides from scrapie fibrils, determining sequences and making such peptides are known in the art. (e.g., Beekes, M., *et al.* J. Gen. Virol. 76: 2567-76, 1995).

For example, one form of GSS has been linked to a PrP mutation at codon 102, while telencephalic GSS segregates with a mutation at codon 117. Mutations at codons

198 and 217 result in <sup>a</sup> form of GSS in which neuritic plaques characteristic of Alzheimer's disease contain PrP instead of A $\beta$  peptide. Certain forms of familial CJD have been associated with mutations at codons 200 and 210; mutations at codons 129 and 178 have been found in both familial CJD and FFI. (Baldwin, *supra*).

5 d. Senile Systemic Amyloidosis

Amyloid deposition, either systemic or focal, increases with age. For example, fibrils of wild type transthyretin (TTR) are commonly found in the heart tissue of elderly individuals. These may be asymptomatic, clinically silent, or may result in heart failure. Asymptomatic fibrillar focal deposits may also occur in the brain (A $\beta$ ), corpora amylacea  
10 of the prostate (A $\beta_2$  microglobulin), joints and seminal vesicles.

e. Cerebral Amyloidosis

Local deposition of amyloid is common in the brain, particularly in elderly individuals. The most frequent type of amyloid in the brain is composed primarily of A $\beta$  peptide fibrils, resulting in dementia or sporadic (non-hereditary) Alzheimer's disease. In  
15 fact, the incidence of sporadic Alzheimer's disease greatly exceeds forms shown to be hereditary. Fibril peptides forming these plaques are very similar to those described above, with reference to hereditary forms of Alzheimer's disease (AD).

f. Dialysis-related Amyloidosis

Plaques composed of  $\beta_2$  microglobulin (A $\beta_2$ M) fibrils commonly develop in  
20 patients receiving long term hemodialysis or peritoneal dialysis.  $\beta_2$  microglobulin is a 11.8 kilodalton polypeptide and is the light chain of Class I MHC antigens, which are present on all nucleated cells. Under normal circumstances, it is continuously shed from cell membranes and is normally filtered by the kidney. Failure of clearance, such as in the case of impaired renal function, leads to deposition in the kidney and other sites (primarily  
25 in collagen-rich tissues of the joints). Unlike other fibril proteins, A $\beta_2$ M molecules are generally present in unfragmented form in the fibrils. (Benson, *supra*).

g. Hormone-derived Amyloidoses

Endocrine organs may harbor amyloid deposits, particularly in aged individuals. Hormone-secreting tumors may also contain hormone-derived amyloid plaques, the fibrils  
30 of which are made up of polypeptide hormones such as calcitonin (medullary carcinoma of the thyroid), islet amyloid polypeptide (amylin; occurring in most patients with Type II

natriuretic

diabetes), and atrial ~~natriuretic~~ peptide (isolated atrial amyloidosis). Sequences and structures of these proteins are well known in the art.

#### h. Miscellaneous Amyloidoses

There are a variety of other forms of amyloid disease that are normally manifest as localized deposits of amyloid. In general, these diseases are probably the result of the localized production and/or lack of catabolism of specific fibril precursors or a predisposition of a particular tissue (such as the joint) for fibril deposition. Examples of such idiopathic deposition include nodular AL amyloid, cutaneous amyloid, endocrine amyloid, and tumor-related amyloid.

10

#### C. Pharmaceutical Compositions

It is the discovery of the present invention that compositions capable of eliciting or providing an immune response directed to certain components of amyloid plaques are effective to treat or prevent development of amyloid diseases. In particular, according to the invention provided herein, it is possible to prevent progression of, ameliorate the symptoms of, and/or reduce amyloid plaque burden in afflicted individuals, when an immunostimulatory dose of an anti-amyloid agent, or corresponding anti-amyloid immune reagent, is administered to the patient. This section describes exemplary anti-amyloid agents that produce active, as well as passive, immune responses to amyloid plaques and provides exemplary data showing the effect treatment using such compositions on amyloid plaque burden.

Generally, anti-amyloid agents of the invention are composed of a specific plaque component, preferably a fibril forming component, which is usually a characteristic protein, peptide, or fragment thereof, as described in the previous section and exemplified below. More generally, therapeutic agents for use in the present invention produce or induce an immune response against a plaque, or more specifically, a fibril component thereof. Such agents therefore include, but are not limited to, the component itself and variants thereof, analogs and mimetics of the component that induce and/or cross-react with antibodies to the component, as well as antibodies or T-cells that are specifically reactive with the amyloid component. According to an important feature, pharmaceutical compositions are not selected from non-specific components - that is, from those components that are generally circulating or that are ubiquitous throughout the body. By way of example, Serum Amyloid Protein (SAP) is a circulating plasma glycoprotein that is

and amyloid  
 produced in the liver ~~and~~ binds to most known forms of amyloid deposits. Therapeutic compositions are preferably directed to this component.

Induction of an immune response can be active, as when an immunogen is administered to induce antibodies or T-cells reactive with the component in a patient, or  
 5 passive, as when an antibody is administered that itself binds to the amyloid component in the patient. Exemplary agents for inducing or producing an immune response against amyloid plaques are described in the sections below.

Pharmaceutical compositions of the present invention may include, in addition to the immunogenic agent(s), an effective amount of an adjuvant and/or an excipient.  
 10 Pharmaceutically effective an useful adjuvants and excipients are well known in the art, and are described in more detail in the Sections that follow.

# 1. Immunostimulatory Agents (Active Immune Response)

## a. Anti-fibril Compositions

15 One general class of preferred anti-amyloid agents consists of agents that are derived from amyloid fibril proteins. As mentioned above, the hallmark of amyloid diseases is the deposition in an organ or organ of amyloid plaques consisting mainly of fibrils, which, in turn, are composed of characteristic fibril proteins or peptides. According to the present invention, such a fibril protein or peptide component is a useful  
 20 agent for inducing an anti-amyloid immune response.

Tables 1 and 2 summarize exemplary fibril-forming proteins that are characteristic of various amyloid diseases. In accordance with this aspect of the present invention, administration to an afflicted or susceptible individual of an immunostimulatory composition which includes the appropriate fibril protein or peptide, including homologs  
 25 or fragments thereof, provides therapeutic or prophylaxis with respect to the amyloid disease.

By way of example, A $\beta$ , also known as  $\beta$ -amyloid peptide, or A4 peptide (see US Patent 4,666,829; Glenner & Wong, Biochem. Biophys. Res. Commun. 120, 1131 (1984)), is a peptide of 39-43 amino acids, which is the principal component of characteristic  
 30 plaques of Alzheimer's disease. A $\beta$  is generated by processing of a larger protein APP by two enzymes, termed  $\beta$  and gamma secretases (see Hardy, TINS 20, 154 (1997)).



describes

Example I describes the results of experiments carried out in support of the present invention, in which A $\beta$ 42 peptide was administered to heterozygote transgenic mice that overexpress human APP with a mutation at position 717. These mice, known as "PDAPP mice" exhibit Alzheimer's-like pathology and are considered to be an animal model for

5 Alzheimer's disease (Games, *et al.*, *Nature* 373: 523-7, 1995). As detailed in the Example, these mice exhibit detectable A $\beta$  plaque neuropathology in their brains beginning at about 6 months of age, with plaque deposition progressing over time. In the experiments described herein, aggregated A $\beta$ 42 (AN1792) was administered to the mice. Most of the treated mice (7/9) had no detectable amyloid in their brains at 13 months of

10 age, in contrast to control mice (saline-injected or untreated), all of which showed significant brain amyloid burden at this age (FIG. 2). These differences were even more pronounced in the hippocampus (FIG. 3). Treated mice also exhibited significant serum antibody titers against A $\beta$  (all greater than 1:1000, 8/9 greater than 1/10,000; FIG. 1, Table 3A). Generally, saline-treated mice exhibited less than 4-5 times background levels of

15 antibodies against A $\beta$  at a dilution of 1:100 at all times tested, and were therefore deemed to have no significant response relative to control (Table 3B). These studies demonstrated that injection with the specific fibril forming peptide A $\beta$  provides protection against deposition of A $\beta$  amyloid plaques.

Serum Amyloid Protein (SAP), is a circulating plasma glycoprotein that is

20 produced in the liver and binds in a calcium-dependent manner to all forms of amyloid fibril, including fibrils of cerebral amyloid plaques in Alzheimer's disease. As part of the foregoing experiments, a group of mice was injected with SAP; these mice developed significant serum titers to SAP (1:1000-1:30000), but did not develop detectable serum titers to A $\beta$  peptide and developed cerebral plaque neuropathology (FIG. 2).

25 Further experiments, detailed in Example II, demonstrate dose dependence of the immunogenic effect of A $\beta$  injections in mice treated between 5 weeks and about 8 months of age. In these mice, mean serum titers of anti-A $\beta$  peptide antibodies increased with the number of immunizations and with increasing dosages; however, after four immunizations, serum titers measured five days following the immunization leveled off

30 over the higher doses (1-300  $\mu$ g) at levels around 1:10000 (FIG. 5).

Additional experiments in support of the present invention are described in Example III, in which PDAPP model mice were treated with A $\beta$ 42 commencing at a time

point (about 11 <sup>months</sup> ~~month~~ of age) after amyloid plaques were already present in their brains. In these studies, the animals were immunized with A $\beta$ 42 or saline, and were sacrificed for amyloid burden testing at age 15 or 18 months. As illustrated in FIG. 7, at 18 months of age, A $\beta$ 42-treated mice exhibited a significantly lower mean amyloid plaque burden (plaque burden, 0.01%) than either PBS-treated 18-month old controls (plaque burden, 4.7%) or 12 month untreated animals (0.28%), where plaque burden is measured by image analysis, as detailed in Example XIII, part 8. These experiments demonstrate the efficacy of the treatment methods of the present invention in reducing existing plaque burden and preventing progression of plaque burden in diseased individuals.

According to this aspect of the invention, therapeutic agents are derived from fibril peptides or proteins which comprise the plaques that are characteristic of the disease of interest. Alternatively, such agents are antigenically similar enough to such components to induce an immune response that also cross-reacts with the fibril component. Tables 1 and 2 provide examples of such fibril peptides and proteins, the compositions and sequences of which are known in the art or can be easily determined according to methods known in the art. (See references cited below and in Section B2 for references that specifically teach methods for extraction and/or compositions of various fibril peptide components; further exemplary fibril components are described below.) Thus, in accordance with the present invention, where a diagnosis of an amyloid disease is made, based on clinical and/or biopsy determinations, the skilled practitioner will be able to ascertain the fibril composition of the amyloid deposits and provide an agent that induces an immune response directed to the fibrillar peptides or proteins.

By way of example, as described above, the therapeutic agent used in treating Alzheimer's disease or other amyloid diseases characterized by A $\beta$  fibril deposition can be any of the naturally occurring forms of A $\beta$  peptide, and particularly the human forms (i.e., A $\beta$ 39, A $\beta$ 40, A $\beta$ 41, A $\beta$ 42 or A $\beta$ 43). The sequences of these peptides and their relationship to the APP precursor are known in the art and are well known in the art (e.g., Hardy et al., TINS 20, 155-158 (1997)). For example, A $\beta$ 42 has the sequence:

H2N-Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Ile-Ala-OH. (SEQ ID NO: 1)

and

A $\beta$ 41, A $\beta$ 40 ~~and~~ A $\beta$ 39 differ from A $\beta$ 42 by the omission of Ala, Ala-Ile, and Ala-Ile-Val respectively from the C-terminal end of the peptide. A $\beta$ 43 differs from A $\beta$ 42 by the presence of a threonine residue at the C-terminus. According to a preferred embodiment of the invention, therapeutic agents will induce an immune response against all or a portion of the fibril component of the disease of interest. For example, a preferred A $\beta$  immunogenic composition is an agent that induces an antibody specific to the free N-terminus of A $\beta$ . Such a composition has the advantage that it would not recognize the precursor protein,  $\beta$ -APP, thereby rendering it less likely to produce autoimmunity.

By way of further example, it is appreciated that patients afflicted with diseases characterized by the deposition of AA fibrils, for example, certain chronic inflammatory disorders, chronic local or systemic microbial infections, and malignant neoplasms, as described above, can be treated with AA peptide, a known 8 kilodalton fragment of serum amyloid A protein (ApoSSA). Exemplary AA amyloid disorders include, but are not limited to inflammatory diseases such as rheumatoid arthritis, juvenile chronic arthritis, ankylosing spondylitis, psoriasis, psoriatic arthropathy, Reiter's syndrome, Adult Still's disease, Behçet's syndrome, Crohn's disease, chronic microbial infections such as leprosy, tuberculosis, bronchiectasis, decubitus ulcers, chronic pyelonephritis, osteomyelitis, and Whipple's disease, as well as malignant neoplasms such as Hodgkin's lymphoma, renal carcinoma, carcinomas of gut, lung and urogenital tract, basal cell carcinoma, and hairy cell leukemia.

AA peptide refers to one or more of a heterogeneous group of peptides derived from the N-terminus of precursor protein serum amyloid A (ApoSSA), commencing at residue 1, 2 or 3 of the precursor protein and ending at any point between residues 58 and 84; commonly AA fibrils are composed of residues 1-76 of ApoSSA. Precise structures and compositions can be determined, and appropriate peptides synthesized according to methods well known in the art (Liepnieks, J.J., *et al.* Biochem. Biophys Acta 1270: 81-86, 1995).

By way of further example, fragments derived from the N-terminal region which contain all or part of the variable (V<sub>L</sub>) domain of immunoglobulin light chains (kappa or lambda chain) generally comprise amyloid deposits in mesenchymal tissues, causing peripheral and autonomic neuropathy, carpal tunnel syndrome, macroglossia, restrictive cardiomyopathy, arthropathy of large joints, immune dyscrasias, myelomas, as well as

*Compositions**preferably*

occult dyscrasias. ~~Compositions~~ of the invention will ~~preferably~~ induce an immune response against a portion of the light chain, preferably against a "neoepitope" – an epitope that is formed as a result of fragmentation of the parent molecule – to reduce possible autoimmune effects.

5 Various hereditary amyloid diseases are also amenable to the treatment methods of the present invention. Such diseases are described in Section B.2, above. For example, various forms of familial amyloid polyneuropathy are the result of at least fifty mutant forms of transthyretin (TTR), a 14 kilodalton protein produced by the liver, each characterized by a single amino acid change. While many of these forms of this disease  
10 are distinguishable on the basis of their particular pathologies and/or demographic origins, it is appreciated that therapeutic compositions may also be composed of agents that induce an immune response against more than one form of TTR, such as a mixture of two or more forms of ATTR, including wildtype TTR, to provide a generally useful therapeutic composition.

15 AapoAI-containing amyloid deposits are found in persons having point mutations in the molecule apolipoprotein AI. Patients with this form of disease generally present with peripheral neuropathy or renal failure. According to the present invention, therapeutic compositions are made up one or more of the various forms of AapoAI described herein or known in the art.

20 Certain familial forms of Alzheimer's disease, as well as Down's syndrome, are the result of mutations in beta amyloid precursor protein, resulting in deposition of plaques having fibrils composed mainly of  $\beta$ -amyloid peptide ( $A\beta$ ). The use of  $A\beta$  peptide in therapeutic compositions of the present invention is described above and exemplified herein.

25 Other formulations for treating hereditary forms of amyloidosis, discussed above, include compositions that produce immune responses against gelsolin fragments for treatment of hereditary systemic amyloidosis, mutant lysozyme protein (Alys), for treatment of a hereditary neuropathy, mutant alpha chain of fibrinogen (AfibA) for a non-neuropathic form of amyloidosis manifest as renal disease, mutant cystatin C (Acys) for  
30 treatment of a form of hereditary cerebral angiopathy reported in Iceland. In addition, certain hereditary forms of prion disease (e.g., Creutzfeldt-Jacob disease (CJD), Gerstmann-Sträussler-Scheinker syndrome (GSS), and fatal familial insomnia (FFI)) are characterized by a mutant isoform of prion protein, PrP<sup>Sc</sup>. This protein can be used in

compositions

deposition

therapeutic compositions for treatment and prevention of deposition of PrP plaques, in accordance with the present invention.

As discussed above, amyloid deposition, either systemic or focal, is also associated with aging. It is a further aspect of the present invention that such deposition can be prevented or treated by administering to susceptible individuals compositions consisting of one or more proteins associated with such aging. Thus, plaques composed of ATTR derived from wild type TTR are frequently found in heart tissue of the elderly. Similarly, certain elderly individuals may develop asymptomatic fibrillar focal deposits of A $\beta$  in their brains; A $\beta$  peptide treatment, as detailed herein may be warranted in such individuals.  $\beta_2$  microglobulin is a frequent component of corpora amylacea of the prostate, and is therefore a further candidate agent in accordance with the present invention.

By way of further example, but not limitation, there are a number of additional, non-hereditary forms amyloid disease that are candidates for treatment methods of the present invention.  $\beta_2$  microglobulin fibrillar plaques commonly develop in patients receiving long term hemodialysis or peritoneal dialysis. Such patients may be treated by treatment with therapeutic compositions directed to  $\beta_2$  microglobulin or, more preferably, immunogenic epitopes thereof, in accordance with the present invention.

Hormone-secreting tumors may also contain hormone-derived amyloid plaques, the composition of which are generally characteristic of the particular endocrine organ affected. Thus such fibrils may be made up of polypeptide hormones such as calcitonin (medullary carcinoma of the thyroid), islet amyloid polypeptide (occurring in most patients with Type II diabetes), and atrial natriuretic peptide (isolated atrial amyloidosis). Compositions directed at amyloid deposits which form in the aortic intima in atherosclerosis are also contemplated by the present invention. For example, Westermarck, *et al.* describe a 69 amino acid N-terminal fragment of Apolipoprotein A which forms such plaques (Westermarck, *et al.* Am. J. Path. 147: 1186-92, 1995); therapeutic compositions of the present invention include immunological reagents directed to such a fragment, as well as the fragment itself.

The foregoing discussion has focused on amyloid fibril components that may be used as therapeutic agents in treating or preventing various forms of amyloid disease. The therapeutic agent can also be an active fragment or analog of a naturally occurring or mutant fibril peptide or protein that contains an epitope that induces a similar protective or

## response

- therapeutic immune ~~response~~ on administration to a human. Immunogenic fragments typically have a sequence of at least 3, 5, 6, 10 or 20 contiguous amino acids from a natural peptide. Exemplary A $\beta$  peptide immunogenic fragments include A $\beta$ 1-5, 1-6, 1-7, 1-10, 3-7, 1-3, 1-4, 1-12, 13-28, 17-28, 1-28, 25-35, 35-40 and 35-42. Fragments
- 5 lacking at least one, and sometimes at least 5 or 10 C-terminal amino acid present in a naturally occurring forms of the fibril component are used in some methods. For example, a fragment lacking 5 amino acids from the C-terminal end of A $\beta$ 43 includes the first 38 amino acids from the N-terminal end of AB. Fragments from the N-terminal half of A $\beta$  are preferred in some methods. Analogs include allelic, species and induced variants.
- 10 Analogs typically differ from naturally occurring peptides at one or a few positions, often by virtue of conservative substitutions. Analogs typically exhibit at least 80 or 90% sequence identity with natural peptides. Some analogs also include unnatural amino acids or modifications of N or C terminal amino acids. Examples of unnatural amino acids are  $\alpha$ ,  $\alpha$ -disubstituted amino acids, N-alkyl amino acids, lactic acid, 4-hydroxyproline,  $\gamma$ -
- 15 carboxyglutamate,  $\gamma$ -N,N,N-trimethyllysine,  $\gamma$ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine,  $\omega$ -N-methylarginine.

- Generally, persons skilled in the art will appreciate that fragments and analogs designed in accordance with this aspect of the invention can be screened for cross-
- 20 reactivity with the naturally occurring fibril components and/or prophylactic or therapeutic efficacy in transgenic animal models as described below. Such fragments or analogs may be used in therapeutic compositions of the present invention, if their immunoreactivity and animal model efficacy is roughly equivalent to or greater than the corresponding parameters measured for the amyloid fibril components.

- 25 Such peptides, proteins, or fragments, analogs and other amyloidogenic peptides can be synthesized by solid phase peptide synthesis or recombinant expression, according to standard methods well known in the art, or can be obtained from natural sources. Exemplary fibril compositions, methods of extraction of fibrils, sequences of fibril peptide or protein components are provided by many of the references cited in conjunction with
- 30 the descriptions of the specific fibril components provided herein. Additionally, other compositions, methods of extracting and determining sequences are known in the art available to persons desiring to make and use such compositions. Automatic peptide

similarity

mimetics

acid sequence ~~similarity~~ with A $\beta$  but nevertheless serve as ~~mimetics~~ of A $\beta$  and induce a similar immune response. For example, any peptides and proteins forming  $\beta$ -pleated sheets can be screened for suitability. Anti-idiotypic antibodies against monoclonal antibodies to A $\beta$  or other amyloidogenic peptides can also be used. Such anti-Id

5 antibodies mimic the antigen and generate an immune response to it (see *Essential Immunology* (Roit ed., Blackwell Scientific Publications, Palo Alto, 6th ed.), p. 181). Agents other than A $\beta$  peptides should induce an immunogenic response against one or more of the preferred segments of A $\beta$  listed above (e.g., 1-10, 1-7, 1-3, and 3-7). Preferably, such agents induce an immunogenic response that is specifically directed to

10 one of these segments without being directed to other segments of A $\beta$ .

Random libraries of peptides or other compounds can also be screened for suitability. Combinatorial libraries can be produced for many types of compounds that can be synthesized in a step-by-step fashion. Such compounds include polypeptides, beta-turn mimetics, polysaccharides, phospholipids, hormones, prostaglandins, steroids, aromatic

15 compounds, heterocyclic compounds, benzodiazepines, oligomeric N-substituted glycines and oligocarbamates. Large combinatorial libraries of the compounds can be constructed by the encoded synthetic libraries (ESL) method described in Affymax, WO 95/12608, Affymax, WO 93/06121, Columbia University, WO 94/08051, Pharmacopeia, WO 95/35503 and Scripps, WO 95/30642 (each of which is incorporated by reference for all

20 purposes). Peptide libraries can also be generated by phage display methods. See, e.g., Devlin, WO 91/18980.

Combinatorial libraries and other compounds are initially screened for suitability by determining their capacity to bind to antibodies or lymphocytes (B or T) known to be specific for A $\beta$  or other amyloidogenic peptides such as ATTR. For example, initial

25 screens can be performed with any polyclonal sera or monoclonal antibody to A $\beta$  or any other amyloidogenic peptide of interest. Compounds identified by such screens are then further analyzed for capacity to induce antibodies or reactive lymphocytes to A $\beta$  or other amyloidogenic peptide. For example, multiple dilutions of sera can be tested on microtiter plates that have been precoated with fibril peptide, and a standard ELISA can be

30 performed to test for reactive antibodies to A $\beta$ . Compounds can then be tested for prophylactic and therapeutic efficacy in transgenic animals predisposed to an amyloidogenic disease, as described in the Examples. Such animals include, for example,

<sup>mutation</sup>  
 mice bearing a 717 ~~mutation~~<sup>supra</sup> of APP described by Games et al., ~~supra~~, and mice bearing a  
 670/671 Swedish mutation of APP such as described by McConlogue et al., US 5,612,486  
 and Hsiao et al., Science 274, 99 (1996); Stausenbiel et al., Proc. Natl. Acad. Sci. USA 94,  
 13287-13292 (1997); Sturchler-Pierrat et al., Proc. Natl. Acad. Sci. USA 94, 13287-13292  
 5 (1997); Borchelt et al., Neuron 19, 939-945 (1997)). The same screening approach can be  
 used on other potential agents such as fragments of A $\beta$ , analogs of A $\beta$  and longer peptides  
 including A $\beta$ , described above.

b. Other Plaque Components

It is appreciated that immunological responses directed at other amyloid plaque  
 10 components can also be effective in preventing, retarding or reducing plaque deposition in  
 amyloid diseases. Such components may be minor components of fibrils or associated  
 with fibrils or fibril formation in the plaques, with the caveat that components that are  
 ubiquitous throughout the body, or relatively non-specific to the amyloid deposit, are  
 generally less suitable for use as therapeutic targets.

It is therefore a further discovery of the present invention that agents that induce an  
 immune response to specific plaque components are useful in treating or preventing  
 progression of amyloid diseases. This section provides background on several exemplary  
 amyloid plaque-associated molecules. Induction of an immune response against any of  
 these molecules, alone or in combination with immunogenic therapeutic compositions  
 20 against the fibril components described above or against any of the other non-fibril  
 forming components described below, provides an additional anti-amyloid treatment  
 regimen, in accordance with the present invention. Also forming part of the present  
 invention are passive immunization regimens based on such plaque components, as  
 described herein.

By way of example, synuclein is a protein that is structurally similar to  
 apolipoproteins but is found in neuronal cytosol, particularly in the vicinity of presynaptic  
 terminals. There are at least three forms of the protein, termed  $\alpha$ ,  $\beta$  and  $\gamma$  synuclein.  
 Recently, it has been shown that  $\alpha$  and  $\beta$  synuclein are involved in nucleation of amyloid  
 deposits in certain amyloid diseases, particularly Alzheimer's disease. (Clayton, D.F., *et*  
 30 *al.*, *TINS* 21(6): 249-255, 1998). More specifically, a fragment of the NAC domain of  $\alpha$   
 and  $\beta$  synuclein (residues 61-95) has been isolated from amyloid plaques in Alzheimer's  
 patients; in fact this fragment comprises about 10% of the plaque that remains insoluble



*with* *George*

after solubilization ~~with~~ sodium dodecyl sulfate (SDS). (~~George~~<sup>George</sup>, J.M., *et al.* Neurosci. News 1: 12-17, 1995). Further, both the full length  $\alpha$  synuclein and the NAC fragment thereof have been reported to accelerate the aggregation of  $\beta$ -amyloid peptide into insoluble amyloid *in vitro*. (Clayton, *supra*).

5 Additional components associated with amyloid plaques include non-peptide components. For example, perlecan and perlecan-derived glycosaminoglycans are large heparin sulfate proteoglycans that are present in A $\beta$ -containing amyloid plaques of Alzheimer's disease and other CNS and systemic amyloidoses, including amylin plaques associated with diabetes. These compounds have been shown to enhance A $\beta$  fibril

10 formation. Both the core protein and glycosaminoglycan chains of perlecan have been shown to participate in binding to A $\beta$ . Additional glycosaminoglycans, specifically, dermatan sulfate, chondroitin-4-sulfate, and pentosan polysulfate, are commonly found in amyloid plaques of various types and have also been shown to enhance fibril formation. Dextran sulfate also has this property. This enhancement is significantly reduced when the

15 molecules are de-sulfated. Immunogenic therapeutics directed against the sulfated forms of glycosaminoglycans, including the specific glycosaminoglycans themselves, form an additional embodiment of the present invention, either as a primary or secondary treatment. Production of such molecules, as well as appropriate therapeutic compositions containing such molecules, is within the skill of the ordinary practitioner in the art.

20

## 2. Agents Inducing Passive Immune Response

Therapeutic agents of the invention also include immune reagents, such as antibodies, that specifically bind to fibril peptides or other components of amyloid plaques. Such antibodies can be monoclonal or polyclonal, and have binding specificities

25 that are consonant with the type of amyloid disease to be targeted. Therapeutic compositions and treatment regimens may include a antibodies directed to a single binding domain or epitope on a particular fibril or non-fibril component of a plaque, or may include antibodies directed to two or more epitopes on the same component or antibodies directed to epitopes on multiple components of the plaque.

30

For example, in experiments carried out in support of the present invention, 8½ to 10½ month old PDAPP mice were given intraperitoneal (i.p.) injections of polyclonal anti-A $\beta$ 42 or monoclonal anti-A $\beta$  antibodies prepared against specific epitopes

of A $\beta$  peptide, or <sup>saline</sup>~~saline~~, as detailed in Example XI herein. In <sup>these</sup>~~these~~ experiments, circulating antibody concentrations were monitored, and booster injections were given as needed to maintain a circulating antibody concentration of greater than 1:1000 with respect to the specific antigen to which the antibody was made. Reductions in total A $\beta$  levels were observed, compared to control, in the cortex, hippocampus and cerebellum brain regions of antibody-treated mice; highest reductions were exhibited in mice treated with polyclonal antibodies in these studies.

In further experiments carried out in support of the invention, a predictive *ex vivo* assay (Example XIV) was used to test clearing of an antibody against a fragment of synuclein referred to as NAC. Synuclein has been shown to be an amyloid plaque-associated protein. An antibody to NAC was contacted with a brain tissue sample containing amyloid plaques and microglial cells. Rabbit serum was used as a control. Subsequent monitoring showed a marked reduction in the number and size of plaques indicative of clearing activity of the antibody.

From these data, it is apparent that amyloid plaque load associated with Alzheimer's disease and other amyloid diseases can be greatly diminished by administration of immune reagents directed against epitopes of A $\beta$  peptide or against the NAC fragment of synuclein, which are effective to reduce amyloid plaque load. It is further understood that a wide variety of antibodies can be used in such compositions. Antibodies that bind specifically to the aggregated form of A $\beta$  without binding to the dissociated form are suitable for use in the invention, as are antibodies that bind specifically to the dissociated form without binding to the aggregated form. Other suitable antibodies bind to both aggregated and dissociated forms. Some such antibodies bind to a naturally occurring short form of A $\beta$  (i.e., A $\beta$ 39, 40 or 41) without binding to a naturally occurring long form of A $\beta$  (i.e., A $\beta$ 42 and A $\beta$ 43). Some antibodies bind to a long form without binding to a short form. Some antibodies bind to A $\beta$  without binding to full-length amyloid precursor protein. Some antibodies bind to A $\beta$  with a binding affinity greater than or equal to about  $10^6$ ,  $10^7$ ,  $10^8$ ,  $10^9$ , or  $10^{10}$  M $^{-1}$ .

Polyclonal sera typically contain mixed populations of antibodies binding to several epitopes along the length of A $\beta$ . Monoclonal antibodies bind to a specific epitope within A $\beta$  that can be a conformational or nonconformational epitope. Some monoclonal antibodies bind to an epitope within residues 1-28 of A $\beta$  (with the first N terminal residue

both

1991), or Chothia & Lesk, *J. Mol. Biol.* 196:901-917 (1987); Chothia et al., *Nature* 342:878-883 (1989).

### **b. Production of Non-human Antibodies**

The production of non-human monoclonal antibodies, e.g., murine, guinea pig, rabbit or rat, can be accomplished by, for example, immunizing the animal with a plaque component, such as A $\beta$  or other fibril components. A longer polypeptide comprising A $\beta$  or an immunogenic fragment of A $\beta$  or anti-idiotypic antibodies to an antibody to A $\beta$  can also be used. See e.g., Harlow & Lane, *Antibodies, A Laboratory Manual* (CSHP NY, 1988) (incorporated by reference for all purposes). Such an immunogen can be obtained from a natural source, by peptide synthesis or by recombinant expression. Optionally, the immunogen can be administered fused or otherwise complexed with a carrier protein, as described below. Optionally, the immunogen can be administered with an adjuvant. Several types of adjuvant can be used as described below. Complete Freund's adjuvant followed by incomplete adjuvant is preferred for immunization of laboratory animals. Rabbits or guinea pigs are typically used for making polyclonal antibodies. Mice are typically used for making monoclonal antibodies. Antibodies are screened for specific binding to the immunogen. Optionally, antibodies are further screened for binding to a specific region of the immunogen. For example, in the case of A $\beta$  peptide as immunogen, screening can be accomplished by determining binding of an antibody to a collection of deletion mutants of an A $\beta$  peptide and determining which deletion mutants bind to the antibody. Binding can be assessed, for example, by Western blot or ELISA. The smallest fragment to show specific binding to the antibody defines the epitope of the antibody. Alternatively, epitope specificity can be determined by a competition assay in which a test and reference antibody compete for binding to the component. If the test and reference antibodies compete, then they bind to the same epitope or epitopes sufficiently proximal that binding of one antibody interferes with binding of the other.

### c. Chimeric and Humanized Antibodies

humanized

similar

Chimeric and ~~humanized~~ antibodies have the same or ~~similar~~ binding specificity and affinity as a mouse or other nonhuman antibody that provides the starting material for construction of a chimeric or humanized antibody. Chimeric antibodies are antibodies whose light and heavy chain genes have been constructed, typically by genetic engineering, from immunoglobulin gene segments belonging to different species. For example, the variable (V) segments of the genes from a mouse monoclonal antibody may be joined to human constant (C) segments, such as IgG1 and IgG4. A typical chimeric antibody is thus a hybrid protein consisting of the V or antigen-binding domain from a mouse antibody and the C or effector domain from a human antibody.

Humanized antibodies have variable region framework residues substantially from a human antibody (termed an acceptor antibody) and complementarity determining regions substantially from a mouse-antibody, (referred to as the donor immunoglobulin). See, Queen et al., *Proc. Natl. Acad. Sci. USA* 86:10029-10033 (1989) and WO 90/07861, US 5,693,762, US 5,693,761, US 5,585,089, US 5,530,101 and Winter, US 5,225,539 (incorporated by reference in their entirety for all purposes). The constant region(s), if present, are also substantially or entirely from a human immunoglobulin. The human variable domains are usually chosen from human antibodies whose framework sequences exhibit a high degree of sequence identity with the murine variable region domains from which the CDRs were derived. The heavy and light chain variable region framework residues can be derived from the same or different human antibody sequences. The human antibody sequences can be the sequences of naturally occurring human antibodies or can be consensus sequences of several human antibodies. See Carter et al., WO 92/22653. Certain amino acids from the human variable region framework residues are selected for substitution based on their possible influence on CDR conformation and/or binding to antigen. Investigation of such possible influences is by modeling, examination of the characteristics of the amino acids at particular locations, or empirical observation of the effects of substitution or mutagenesis of particular amino acids.

For example, when an amino acid differs between a murine variable region framework residue and a selected human variable region framework residue, the human framework amino acid should usually be substituted by the equivalent framework amino acid from the mouse antibody when it is reasonably expected that the amino acid:

- (1) noncovalently binds antigen directly,

- is
- (2) ~~is~~ adjacent to a CDR region,
  - (3) otherwise interacts with a CDR region (e.g. is within about 6 Å of a CDR region), or
  - (4) participates in the VL-VH interface.

5 Other candidates for substitution are acceptor human framework amino acids that are unusual for a human immunoglobulin at that position. These amino acids can be substituted with amino acids from the equivalent position of the mouse donor antibody or from the equivalent positions of more typical human immunoglobulins. Other candidates for substitution are acceptor human framework amino acids that are unusual for  
 10 a human immunoglobulin at that position.. The variable region frameworks of humanized immunoglobulins usually show at least 85% sequence identity to a human variable region framework sequence or consensus of such sequences.

#### d. Human Antibodies

15 Human antibodies against A $\beta$  are provided by a variety of techniques described below. Some human antibodies are selected by competitive binding experiments, or otherwise, to have the same epitope specificity as a particular mouse antibody, such as one of the mouse monoclonals described in Example XI. Human antibodies can also be screened for a particular epitope specificity by using only a  
 20 fragment of A $\beta$  as the immunogen, and/or by screening antibodies against a collection of deletion mutants of A $\beta$ .

#### (1) Trioma Methodology

The basic approach and an exemplary cell fusion partner, SPAZ-4, for use in this approach have been described by Oestberg et al., *Hybridoma* 2:361-367 (1983);  
 25 Oestberg, U.S. Patent No. 4,634,664; and Engleman et al., US Patent 4,634,666 (each of which is incorporated by reference in its entirety for all purposes). The antibody-producing cell lines obtained by this method are called triomas, because they are descended from three cells--two human and one mouse. Initially, a mouse myeloma line is fused with a human B-lymphocyte to obtain a non-antibody-producing xenogeneic hybrid  
 30 cell, such as the SPAZ-4 cell line described by Oestberg, *supra*. The xenogeneic cell is then fused with an immunized human B-lymphocyte to obtain an antibody-producing

<sup>Triomas</sup> trioma cell line. <sup>more</sup> ~~Triomas~~ have been found to produce antibody ~~more~~ stably than ordinary hybridomas made from human cells.

The immunized B-lymphocytes are obtained from the blood, spleen, lymph nodes or bone marrow of a human donor. If antibodies against a specific antigen or epitope are desired, it is preferable to use that antigen or epitope thereof for immunization. Immunization can be either in vivo or in vitro. For in vivo immunization, B cells are typically isolated from a human immunized with A $\beta$ , a fragment thereof, larger polypeptide containing A $\beta$  or fragment, or an anti-idiotypic antibody to an antibody to A $\beta$ . In some methods, B cells are isolated from the same patient who is ultimately to be administered antibody therapy. For in vitro immunization, B-lymphocytes are typically exposed to antigen for a period of 7-14 days in a media such as RPMI-1640 (see Engleman, *supra*) supplemented with 10% human plasma.

The immunized B-lymphocytes are fused to a xenogeneic hybrid cell such as SPAZ-4 by well known methods. For example, the cells are treated with 40-50% polyethylene glycol of MW 1000-4000, at about 37 degrees C, for about 5-10 min. Cells are separated from the fusion mixture and propagated in media selective for the desired hybrids (e.g., HAT or AH). Clones secreting antibodies having the required binding specificity are identified by assaying the trioma culture medium for the ability to bind to A $\beta$  or a fragment thereof. Triomas producing human antibodies having the desired specificity are subcloned by the limiting dilution technique and grown in vitro in culture medium. The trioma cell lines obtained are then tested for the ability to bind A $\beta$  or a fragment thereof.

Although triomas are genetically stable they do not produce antibodies at very high levels. Expression levels can be increased by cloning antibody genes from the trioma into one or more expression vectors, and transforming the vector into standard mammalian, bacterial or yeast cell lines, according to methods well known in the art.

## (2) Transgenic Non-human Mammals

Human antibodies against A $\beta$  can also be produced from non-human transgenic mammals having transgenes encoding at least a segment of the human immunoglobulin locus. Usually, the endogenous immunoglobulin locus of such transgenic mammals is functionally inactivated. Preferably, the segment of the human

combination

See

more efficient in ~~combination~~ with phage-display technology. ~~See~~, e.g., Dower et al., WO 91/17271 and McCafferty et al., WO 92/01047, US 5,877,218, US 5,871,907, US 5,858,657, US 5,837,242, US 5,733,743 and US 5,565,332 (each of which is incorporated by reference in its entirety for all purposes). In these methods, libraries of phage are  
 5 produced in which members display different antibodies on their outer surfaces. Antibodies are usually displayed as Fv or Fab fragments. Phage displaying antibodies with a desired specificity are selected by affinity enrichment to an A $\beta$  peptide or fragment thereof.

In a variation of the phage-display method, human antibodies having the binding  
 10 specificity of a selected murine antibody can be produced. See Winter, WO 92/20791. In this method, either the heavy or light chain variable region of the selected murine antibody is used as a starting material. If, for example, a light chain variable region is selected as the starting material, a phage library is constructed in which members display the same light chain variable region (i.e., the murine starting material) and a different heavy chain  
 15 variable region. The heavy chain variable regions are obtained from a library of rearranged human heavy chain variable regions. A phage showing strong specific binding for the component of interest (e.g., at least  $10^8$  and preferably at least  $10^9$  M<sup>-1</sup>) is selected. The human heavy chain variable region from this phage then serves as a starting material for constructing a further phage library. In this library, each phage displays the same  
 20 heavy chain variable region (i.e., the region identified from the first display library) and a different light chain variable region. The light chain variable regions are obtained from a library of rearranged human variable light chain regions. Again, phage showing strong specific binding for amyloid peptide component are selected. These phage display the variable regions of completely human anti-amyloid peptide antibodies. These antibodies  
 25 usually have the same or similar epitope specificity as the murine starting material.

#### e. Selection of Constant Region

The heavy and light chain variable regions of chimeric, humanized, or human antibodies can be linked to at least a portion of a human constant region. The choice of constant region depends, in part, whether antibody-dependent complement  
 30 and/or cellular mediated toxicity is desired. For example, isotopes IgG1 and IgG3 have complement activity and isotopes IgG2 and IgG4 do not. Choice of isotype can also affect passage of antibody into the brain. Light chain constant regions can be lambda or kappa. Antibodies can be expressed as tetramers containing two light and two heavy

*heavy* *(ab)2*

chains, as separate ~~heavy~~ chains, light chains, as Fab, Fab' ~~F(ab)2~~, and Fv, or as single chain antibodies in which heavy and light chain variable domains are linked through a spacer.

#### 5 f. Expression of Recombinant Antibodies

Chimeric, humanized and human antibodies are typically produced by recombinant expression. Recombinant polynucleotide constructs typically include an expression control sequence operably linked to the coding sequences of antibody chains, including naturally-associated or heterologous promoter regions. Preferably, the

10 expression control sequences are eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells. Once the vector has been incorporated into the appropriate host, the host is maintained under conditions suitable for high level expression of the nucleotide sequences, and the collection and purification of the crossreacting antibodies.

15 These expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA. Commonly, expression vectors contain selection markers, e.g., ampicillin-resistance or hygromycin-resistance, to permit detection of those cells transformed with the desired DNA sequences.

*E. coli* is one prokaryotic host particularly useful for cloning the DNA

20 sequences of the present invention. Microbes, such as yeast are also useful for expression. *Saccharomyces* is a preferred yeast host, with suitable vectors having expression control sequences, an origin of replication, termination sequences and the like as desired. Typical promoters include 3-phosphoglycerate kinase and other glycolytic enzymes. Inducible yeast promoters include, among others, promoters from alcohol dehydrogenase,

25 isocytochrome C, and enzymes responsible for maltose and galactose utilization.

Mammalian cells are a preferred host for expressing nucleotide segments encoding immunoglobulins or fragments thereof. See Winnacker, From Genes to Clones, (VCH Publishers, NY, 1987). A number of suitable host cell lines capable of secreting intact heterologous proteins have been developed in the art, and include CHO cell lines,

30 various COS cell lines, HeLa cells, L cells and myeloma cell lines. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter, an enhancer (Queen et al., *Immunol. Rev.* 89:49 (1986)), and necessary processing information sites, such as ribosome binding sites, RNA splice sites,



peptide 83

activated against the peptide. See Peterson et al., US 5,314,815. Insect cell lines expressing an MHC class II antigen can similarly be used to activate CD4 T cells.

#### 5. Carrier Proteins

5           Some agents for inducing an immune response contain the appropriate epitope for inducing an immune response against amyloid deposits but are too small to be immunogenic. In this situation, a peptide immunogen can be linked to a suitable carrier to help elicit an immune response. Suitable carriers include serum albumins, keyhole limpet hemocyanin, immunoglobulin molecules, thyroglobulin, ovalbumin, tetanus toxoid, or a  
10   toxoid from other pathogenic bacteria, such as diphtheria, *E. coli*, cholera, or *H. pylori*, or an attenuated toxin derivative. Other carriers include T-cell epitopes that bind to multiple MHC alleles, e.g., at least 75% of all human MHC alleles. Such carriers are sometimes known in the art as "universal T-cell epitopes." Examples of universal T-cell epitopes include:

- 15   Influenza Hemagglutinin: HA<sub>307-319</sub> PKYVKQNTLKLAT (SEQ ID NO: 1)  
 PADRE (common residues bolded) AKXVAAWTLKAAA (SEQ ID NO: 2)  
 Malaria CS: T3 epitope EKKIAMEKASSVFNV (SEQ ID NO: 3)  
 Hepatitis B surface antigen: HBsAg<sub>19-28</sub> FFLLTRILTI (SEQ ID NO: 4)  
 Heat Shock Protein 65: hsp65<sub>153-171</sub> DQSIGDLIAEAMDKVGNEG (SEQ ID NO: 5)  
 20   bacille Calmette-Guerin QVHFQPLPPAVVKL (SEQ ID NO: 6)  
 Tetanus toxoid: TT<sub>830-844</sub> QYIKANSKFIGITEL (SEQ ID NO: 7)  
 Tetanus toxoid: TT<sub>947-967</sub> FNNFTVSFWLRVPKVSASHLE (SEQ ID NO: 8)  
 HIV gp120 T1: KQIINMWQEVGKAMYA. (SEQ ID NO: 9)

25           Other carriers for stimulating or enhancing an immune response include cytokines such as IL-1, IL-1  $\alpha$  and  $\beta$  peptides, IL-2,  $\gamma$ INF, IL-10, GM-CSF, and chemokines, such as MIP1 $\alpha$  and  $\beta$  and RANTES. Immunogenic agents can also be linked to peptides that enhance transport across tissues, as described in O'Mahony, WO 97/17613 and WO 97/17614.

30           Immunogenic agents can be linked to carriers by chemical crosslinking. Techniques for linking an immunogen to a carrier include the formation of disulfide linkages using N-succinimidyl-3-(2-pyridyl-thio) propionate (SPDP) and succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC) (if the peptide lacks a  
 35   sulfhydryl group, this can be provided by addition of a cysteine residue). These reagents create a disulfide linkage between themselves and peptide cysteine residues on one protein

and an amide linkage <sup>through</sup> ~~through~~ the  $\epsilon$ -amino on a lysine, or <sup>other free</sup> ~~other~~ amino group in other amino acids. A variety of such disulfide/amide-forming agents are described by *Immun. Rev.* 62, 185 (1982). Other bifunctional coupling agents form a thioether rather than a disulfide linkage. Many of these thio-ether-forming agents are commercially available and include reactive esters of 6-maleimidocaproic acid, 2-bromoacetic acid, and 2-iodoacetic acid, 4-(N-maleimido-methyl)cyclohexane-1-carboxylic acid. The carboxyl groups can be activated by combining them with succinimide or 1-hydroxyl-2-nitro-4-sulfonic acid, sodium salt.

Immunogenic peptides can also be expressed as fusion proteins with carriers (i.e., heterologous peptides). The immunogenic peptide can be linked at its amino terminus, its carboxyl terminus, or both to a carrier. Optionally, multiple repeats of the immunogenic peptide can be present in the fusion protein. Optionally, an immunogenic peptide can be linked to multiple copies of a heterologous peptide, for example, at both the N and C termini of the peptide. Some carrier peptides serve to induce a helper T-cell response against the carrier peptide. The induced helper T-cells in turn induce a B-cell response against the immunogenic peptide linked to the carrier peptide.


Some agents of the invention comprise a fusion protein in which an N-terminal fragment of A $\beta$  is linked at its C-terminus to a carrier peptide. In such agents, the N-terminal residue of the fragment of A $\beta$  constitutes the N-terminal residue of the fusion protein. Accordingly, such fusion proteins are effective in inducing antibodies that bind to an epitope that requires the N-terminal residue of A $\beta$  to be in free form. Some agents of the invention comprises a plurality of repeats of an N-terminal segment of A $\beta$  linked at the C-terminus to one or more copy of a carrier peptide. The N-terminal fragment of A $\beta$  incorporated into such fusion proteins sometimes begins at A $\beta$ 1-3 and ends at A $\beta$ 7-11. A $\beta$ 1-7, A $\beta$ 1-3, 1-4, 1-5, and 3-7 are preferred N-terminal fragment of A $\beta$ . Some fusion proteins comprise different N-terminal segments of A $\beta$  in tandem. For example, a fusion protein can comprise A $\beta$ 1-7 followed by A $\beta$ 1-3 followed by a heterologous peptide.

In some fusion proteins, an N-terminal segment of A $\beta$  is fused at its N-terminal end to a heterologous carrier peptide. The same variety of N-terminal segments of A $\beta$  can be used as with C-terminal fusions. Some fusion proteins comprise a heterologous peptide linked to the N-terminus of an N-terminal segment of A $\beta$ , which is in turn linked to one or more additional N-terminal segments of A $\beta$  in tandem.

*examples*                      *use*  
 Some ~~examples~~ of fusion proteins suitable for ~~use~~ in the invention are

shown below. Some of these fusion proteins comprise segments of A $\beta$  linked to tetanus  
 toxoid epitopes such as described in US 5,196,512, EP 378,881 and EP 427,347. Some  
 fusion proteins comprises segments of A $\beta$  linked to carrier peptides described in US  
 5,736,142. Some heterologous peptides are universal T-cell epitopes. In some methods,  
 the agent for administration is simply a single fusion protein with an A $\beta$  segment linked to  
 a heterologous segment in linear configuration. In some methods, the agent is multimer of  
 fusion proteins represented by the formula 2<sup>x</sup>, in which x is an integer from 1-5.  
 Preferably x is 1, 2 or 3, with 2 being most preferred. When x is two, such a multimer has  
 four fusion proteins linked in a preferred configuration referred to as MAP4 (see US  
 5,229,490). Epitopes of A $\beta$  are underlined.

The MAP4 configuration is shown below, where branched structures are  
 produced by initiating peptide synthesis at both the N terminal and side chain amines of  
 lysine. Depending upon the number of times lysine is incorporated into the sequence and  
 allowed to branch, the resulting structure will present multiple N termini. In this example,  
 four identical N termini have been produced on the branched lysine-containing core. Such  
 multiplicity greatly enhances the responsiveness of cognate B cells.



AN90549 (A $\beta$  1-7/Tetanus toxoid 830-844 in a MAP4 configuration):

DAEFRHDQYIKANSKFIGITEL (SEQ ID NO: 10)

AN90550 (A $\beta$  1-7/Tetanus toxoid 947-967 in a MAP4 configuration):

DAEFRHDFNNFTVSFWLRVPKVSASHLE (SEQ ID NO: 11)

AN90542 (A $\beta$  1-7/Tetanus toxoid 830-844 + 947-967 in a linear configuration):

DAEFRHDQYIKANSKFIGITELFNNFTVSFWLRVPKVSASHLE (SEQ ID NO: 12)

AN90576: (A $\beta$  3-9)/Tetanus toxoid 830-844 in a MAP4 configuration):

EFRHDSGQYIKANSKFIGITEL (SEQ ID NO: 13)

US

Peptide described in ~~US~~ 5,736,142 (all in linear configurations):

AN90562 (A $\beta$  1-7/ peptide) AKXVAAWTLKAAADAEFRHD (SEQ ID NO: 14)

AN90543 (A $\beta$  1-7 x 3/ peptide): DAEFRHDDAEFRHDDAEFRHDAKXVAAWTLKAAA  
(SEQ ID NO: 15)

5

Other examples of fusion proteins (immunogenic epitope of A $\beta$  bolded) include  
AKXVAAWTLKAAA-**DAEFRHD-DAEFRHD-DAEFRHD** (SEQ ID

NO: 16)

**DAEFRHD**-AKXVAAWTLKAAA (SEQ ID NO: 17)

10

**DAEFRHD**-ISQAVHAAHAENEAGR (SEQ ID NO: 18)

**FRHDSGY**-ISQAVHAAHAENEAGR (SEQ ID NO: 19)

**EFRHDSG**-ISQAVHAAHAENEAGR (SEQ ID NO: 20)

**PKYVKQNTLKLAT-DAEFRHD-DAEFRHD-DAEFRHD** (SEQ ID

NO: 21)

15

**DAEFRHD-PKYVKQNTLKLAT-DAEFRHD** (SEQ ID NO: 22)

**DAEFRHD-DAEFRHD-DAEFRHD-PKYVKQNTLKLAT** (SEQ ID

NO: 23)

**DAEFRHD-DAEFRHD-PKYVKQNTLKLAT** (SEQ ID NO: 24)

20

**DAEFRHD-PKYVKQNTLKLAT-EKKIAKMEKASSVFNV-**

**QYIKANSKFIGITEL-FNNFTVSFWLRVPKVSASHLE-DAEFRHD**

**DAEFRHD-DAEFRHD-DAEFRHD-QYIKANSKFIGITEL-**

**FNNFTVSFWLRVPKVSASHLE** (SEQ ID

NO: 25)

**DAEFRHD-QYIKANSKFIGITELCFNNFTVSFWLRVPKVSASHLE**

25 (SEQ ID NO: 26)

**DAEFRHD-QYIKANSKFIGITELCFNNFTVSFWLRVPKVSASHLE-**

**DAEFRHD**

(SEQ ID NO: 27)

**DAEFRHD-QYIKANSKFIGITEL** (SEQ ID NO: 28) on a 2

30

branched resin

peptide

Lys-Gly-Cys

peptide

**EQVTNVGGAISQAVHAAHAENEAGR** (Synuclein fusion protein in

35 MAP-4 configuration; SEQ ID NO: 29)

The same or similar carrier proteins and methods of linkage can be used for  
generating immunogens to be used in generation of antibodies against A $\beta$  for use in  
passive immunization. For example, A $\beta$  or a fragment linked to a carrier can be  
administered to a laboratory animal in the production of monoclonal antibodies to A $\beta$ .

40

## 6. Nucleic Acid Encoding Therapeutic Agents

responses

also be

Immune ~~responses~~ against amyloid deposits can ~~also be~~ induced by administration of nucleic acids encoding selected peptide immunogens, or antibodies and their component chains used for passive immunization. Such nucleic acids can be DNA or RNA. A nucleic acid segment encoding an immunogen is typically linked to regulatory elements, such as a promoter and enhancer, that allow expression of the DNA segment in the intended target cells of a patient. For expression in blood cells, as is desirable for induction of an immune response, promoter and enhancer elements from light or heavy chain immunoglobulin genes or the CMV major intermediate early promoter and enhancer are suitable to direct expression. The linked regulatory elements and coding sequences are often cloned into a vector. For administration of double-chain antibodies, the two chains can be cloned in the same or separate vectors.

A number of viral vector systems are available including retroviral systems (see, e.g., Lawrie and Tumin, *Cur. Opin. Genet. Develop.* 3, 102-109, 1993); adenoviral vectors (see, e.g., Bett et al., *J. Virol.* 67, 5911, 1993); adeno-associated virus vectors (see, e.g., Zhou et al., *J. Exp. Med.* 179, 1867, 1994), viral vectors from the pox family including vaccinia virus and the avian pox viruses, viral vectors from the alpha virus genus such as those derived from Sindbis and Semliki Forest Viruses (see, e.g., Dubensky et al., *J. Virol.* 70, 508-519, 1996), Venezuelan equine encephalitis virus (see US 5,643,576) and rhabdoviruses, such as vesicular stomatitis virus (see WO 96/34625) and papillomaviruses (Ohe et al., *Human Gene Therapy* 6, 325-333, 1995); Woo et al., WO 94/12629 and Xiao & Brandsma, *Nucleic Acids. Res.* 24, 2630-2622, 1996).

DNA encoding an immunogen, or a vector containing the same, can be packaged into liposomes. Suitable lipids and related analogs are described by US 5,208,036, 5,264,618, 5,279,833 and 5,283,185. Vectors and DNA encoding an immunogen can also be adsorbed to or associated with particulate carriers, examples of which include polymethyl methacrylate polymers and polylactides and poly(lactide-co-glycolides), see, e.g., McGee et al., *J. Micro Encap.* (1996).

Gene therapy vectors or naked DNA can be delivered in vivo by administration to an individual patient, typically by systemic administration (e.g., intravenous, intraperitoneal, intranasal, gastric, intradermal, intramuscular, subdermal, or intracranial infusion) or topical application (see e.g., US 5,399,346). Such vectors can further include facilitating agents such as bupivacaine (US 5,593,970). DNA can also be administered using a gene gun. See Xiao & Brandsma, *supra*. The DNA encoding an immunogen is

*surface*                      *microprojectiles*

precipitated onto the ~~surface~~ of microscopic metal beads. The ~~microprojectiles~~ are accelerated with a shock wave or expanding helium gas, and penetrate tissues to a depth of several cell layers. For example, The Accel™ Gene Delivery Device manufactured by Agracetus, Inc., (Middleton, WI) is suitable. Alternatively, naked DNA can pass through skin into the blood stream simply by spotting the DNA onto skin with chemical or mechanical irritation (see WO 95/05853).

In a further variation, vectors encoding immunogens can be delivered to cells ex vivo, such as cells explanted from an individual patient (e.g., lymphocytes, bone marrow aspirates, tissue biopsy) or universal donor hematopoietic stem cells, followed by reimplantation of the cells into a patient, usually after selection for cells which have incorporated the vector.

#### 7. Screening Antibodies for Clearing Activity

Example XIV describes methods of screening an antibody for activity in clearing an amyloid deposit. To screen for activity against an amyloid deposit, a tissue sample from a patient with amyloidosis, such as brain tissue in Alzheimer's disease, or an animal model having characteristic amyloid pathology is contacted with phagocytic cells bearing an Fc receptor, such as microglial cells, and the antibody under test in a medium in vitro. The phagocytic cells can be a primary culture or a cell line, such as BV-2, C8-B4, or THP-1. These components are combined on a microscope slide to facilitate microscopic monitoring, or multiple reactions may be performed in parallel in the wells of a microtiter dish. In such a format, a separate miniature microscope slide can be mounted in the separate wells, or a nonmicroscopic detection format, such as ELISA detection of A $\beta$  can be used. Preferably, a series of measurements is made of the amount of amyloid deposit in the in vitro reaction mixture, starting from a baseline value before the reaction has proceeded, and one or more test values during the reaction. The antigen can be detected by staining, for example, with a fluorescently labelled antibody to A $\beta$  or other component of amyloid plaques. The antibody used for staining may or may not be the same as the antibody being tested for clearing activity. A reduction relative to baseline during the reaction of the amyloid deposits indicates that the antibody under test has clearing activity. Such antibodies are likely to be useful in preventing or treating Alzheimer's and other amyloidogenic diseases. As described above, experiments carried out in support of the present invention revealed, using such an assay, that antibodies to the NAC fragment of

Patients amenable to treatment include individuals at risk of disease but not showing symptoms, as well as patients presently showing symptoms of amyloidosis. In the case of Alzheimer's disease, virtually anyone is at risk of suffering from Alzheimer's disease if he or she lives long enough. Therefore, the present methods can be administered prophylactically to the general population without the need for any assessment of the risk of the subject patient. The present methods are especially useful for individuals who do have a known genetic risk of Alzheimer's disease or any of the other hereditary amyloid diseases. Such individuals include those having relatives who have experienced this disease, and those whose risk is determined by analysis of genetic or biochemical markers. Genetic markers of risk toward Alzheimer's disease include mutations in the APP gene, particularly mutations at position 717 and positions 670 and 671 referred to as the Hardy and Swedish mutations respectively (see Hardy, TINS, supra). Other markers of risk are mutations in the presenilin genes, PS1 and PS2, and ApoE4, family history of AD, hypercholesterolemia or atherosclerosis. Individuals presently suffering from Alzheimer's disease can be recognized from characteristic dementia, as well as the presence of risk factors described above. In addition, a number of diagnostic tests are available for identifying individuals who have AD. These include measurement of CSF tau and A $\beta$ 42 levels. Elevated tau and decreased A $\beta$ 42 levels signify the presence of AD. Individuals suffering from Alzheimer's disease can also be diagnosed by MMSE or ADRDA criteria as discussed in the Examples section.

50

of amyloidosis

Other forms ~~of amyloidosis~~ often go undiagnosed, unless a particular predilection for the disease is suspected. One prime symptom is the presence of cardiac or renal disease in a middle-aged to elderly patient who also has signs of other organ involvement. Low voltage or extreme axis deviations of the electrocardiogram and thickened ventricular tissue may be indicative of cardiac involvement. Proteinuria is a symptom of renal involvement. Hepatic involvement may also be suspected, if hepatomegaly is detected by physical examination of the patient. Peripheral neuropathy is also a common occurrence in certain forms of amyloidoses; autonomic neuropathy, characterized by postural hypotension, may also be found. Amyloidosis should be suspected in anyone with a progressive neuropathy of indeterminate origin. A definitive diagnosis of the disease can be made using tissue biopsy methods, where the affected organ(s) are available. For systemic amyloidoses, a fat pad aspirated or rectal biopsy samples may be used. The biopsy material is stained with Congo red, with positive samples exhibiting apple green birefringence under polarized light microscopy.

#### E. Treatment Regimens

In prophylactic applications, pharmaceutical compositions or medicaments are administered to a patient susceptible to, or otherwise at risk of, a particular disease in an amount sufficient to eliminate or reduce the risk or delay the outset of the disease. In therapeutic applications, compositions or medicaments are administered to a patient suspected of, or already suffering from such a disease in an amount sufficient to cure, or at least partially arrest, the symptoms of the disease and its complications. An amount adequate to accomplish this is defined as a therapeutically- or pharmaceutically-effective dose. In both prophylactic and therapeutic regimes, agents are usually administered in several dosages until a sufficient immune response has been achieved. Typically, the immune response is monitored and repeated dosages are given if the immune response starts to wane.

Effective doses of the compositions of the present invention, for the treatment of the above described conditions vary depending upon many different factors, including means of administration, target site, physiological state of the patient, whether the patient is human or an animal, other medications administered, and whether treatment is prophylactic or therapeutic. Usually, the patient is a human, but in some diseases, such as prion protein-associated mad cow disease, the patient can be a nonhuman mammal, such as a bovine. Treatment dosages need to be titrated to optimize safety and efficacy. The



immunogen depends administered  
 amount of immunogen depends on whether adjuvant is also administered, with higher dosages generally being required in the absence of adjuvant. Depending on the immunogenicity of the particular formulation, an amount of an immunogen for administration may vary from 1  $\mu$ g-500  $\mu$ g per patient and more usually from 5-500  $\mu$ g per injection for human administration. Occasionally, a higher dose of 0.5-5 mg per injection is used. Typically at least about 10, 20, 50 or 100  $\mu$ g is used for each human injection. The timing of injections can vary significantly from once a day, to once a year, to once a decade, with successive "boosts" of immunogen somewhat preferred. Generally, in accordance with the teachings provided herein, effective dosages can be monitored by obtaining a fluid sample from the patient, generally a blood serum sample, and determining the titer of antibody developed against the immunogen, using methods well known in the art and readily adaptable to the specific antigen to be measured. Ideally, a sample is taken prior to initial dosing; subsequent samples are taken and titered after each immunization. Generally, a dose or dosing schedule which provides a detectable titer at least four times greater than control or "background" levels at a serum dilution of 1:100 is desirable, where background is defined relative to a control serum or relative to a plate background in ELISA assays. Titers of at least 1:1000 or 1:5000 are preferred in accordance with the present invention.

On any given day that a dosage of immunogen is given, the dosage is usually greater than about 1  $\mu$ g/patient and preferably greater than 10  $\mu$ g/patient if adjuvant is also administered, and at least greater than 10  $\mu$ g/patient and usually greater than 100  $\mu$ g/patient in the absence of adjuvant. Doses for individual immunogens, selected in accordance with the present invention, are determined according to standard dosing and titering methods, taken in conjunction with the teachings provided herein. A typical regimen consists of an immunization followed by booster injections at time intervals, such as 6 week intervals. Another regimen consists of an immunization followed by booster injections 1, 2 and 12 months later. Another regimen entails an injection every two months for life. Alternatively, booster injections can be on an irregular basis as indicated by monitoring of immune response.

For passive immunization with an antibody, the dosage ranges from about 0.0001 to 100 mg/kg, and more usually 0.01 to 5 mg/kg, of the host body weight. For example dosages can be 1 mg/kg body weight or 10 mg/kg body weight. An exemplary treatment

## administration

regime entails ~~administration~~<sup>a</sup> once per every two weeks or once <sup>a</sup> month or once every 3 to 6 months. In some methods, two or more monoclonal antibodies with different binding specificities are administered simultaneously, in which case the dosage of each antibody administered falls within the ranges indicated. Antibody is usually administered on multiple occasions. Intervals between single dosages can be weekly, monthly or yearly. Intervals can also be irregular as indicated by measuring blood levels of antibody to A $\beta$  in the patient. Alternatively, antibody can be administered as a sustained release formulation, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life of the antibody in the patient. In general, human antibodies show the longest half life, followed by humanized antibodies, chimeric antibodies, and nonhuman antibodies. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or therapeutic. In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest of their lives. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the patient shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

Doses for nucleic acids encoding immunogens range from about 10 ng to 1 g, 100 ng to 100 mg, 1  $\mu$ g to 10 mg, or 30-300  $\mu$ g DNA per patient. Doses for infectious viral vectors vary from 10-100, or more, virions per dose.

Agents for inducing an immune response can be administered by parenteral, topical, intravenous, oral, subcutaneous, intraperitoneal, intranasal or intramuscular means for prophylactic and/or therapeutic treatment. Typical routes of administration of an immunogenic agent are intramuscular (i.m.), intravenous (i.v.) or subcutaneous (s.c.), although other routes can be equally effective. Intramuscular injection is most typically performed in the arm or leg muscles. In some methods, agents are injected directly into a particular tissue where deposits have accumulated, for example intracranial injection. Intramuscular injection or intravenous infusion are preferred for administration of antibody. In some methods, particular therapeutic antibodies are injected directly into the cranium. In some methods, antibodies are administered as a sustained release composition or device, such as a Medipad<sup>TM</sup> device.

invention

in

Agents of the ~~invention~~ can optionally be administered ~~in~~ combination with other agents that are at least partly effective in treatment of amyloidogenic disease. In the case of Alzheimer's and Down's syndrome, in which amyloid deposits occur in the brain, agents of the invention can also be administered in conjunction with other agents that increase

5 passage of the agents of the invention across the blood-brain barrier. Further, therapeutic cocktails comprising immunogens designed to provoke an immune response against more than one amyloid component are also contemplated by the present invention, as are a combination of an antibody directed against one plaque component and an immunogen directed to a different plaque component.

10 Immunogenic agents of the invention, such as peptides, are sometimes administered in combination with an adjuvant. A variety of adjuvants can be used in combination with a peptide, such as A $\beta$ , to elicit an immune response. Preferred adjuvants augment the intrinsic response to an immunogen without causing conformational changes in the immunogen that affect the qualitative form of the response. Preferred adjuvants

15 include aluminum hydroxide and aluminum phosphate, 3 De-O-acylated monophosphoryl lipid A (MPL™) (see GB 2220211 (RIBI ImmunoChem Research Inc., Hamilton, Montana, now part of Corixa). Stimulon™ QS-21 is a triterpene glycoside or saponin isolated from the bark of the Quillaja Saponaria Molina tree found in South America (see Kensil et al., in *Vaccine Design: The Subunit and Adjuvant Approach* (eds. Powell &

20 Newman, Plenum Press, NY, 1995); US Patent No. 5,057,540), (Aquila BioPharmaceuticals, Framingham, MA). Other adjuvants are oil in water emulsions (such as squalene or peanut oil), optionally in combination with immune stimulants, such as monophosphoryl lipid A (see Stoute et al., *N. Engl. J. Med.* 336, 86-91 (1997)). Another adjuvant is CpG (WO 98/40100). Alternatively, A $\beta$  can be coupled to an adjuvant.

25 However, such coupling should not substantially change the conformation of A $\beta$  so as to affect the nature of the immune response thereto. Adjuvants can be administered as a component of a therapeutic composition with an active agent or can be administered separately, before, concurrently with, or after administration of the therapeutic agent.

A preferred class of adjuvants is aluminum salts (alum), such as aluminum

30 hydroxide, aluminum phosphate, aluminum sulfate. Such adjuvants can be used with or without other specific immunostimulating agents such as MPL or 3-DMP, QS-21, polymeric or monomeric amino acids such as polyglutamic acid or polylysine. Another

oil-in-water

adjuvants

- class of adjuvants is ~~oil-in-water~~ emulsion formulations. Such ~~adjuvants~~ can be used with or without other specific immunostimulating agents such as muramyl peptides (e.g., N-acetylmuramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-
- 5 2'dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), N-acetylglucosaminyl-N-acetylmuramyl-L-Al-D-isoglu-L-Ala-dipalmitoxy propylamide (DTP-DPP) theramide™, or other bacterial cell wall components. Oil-in-water emulsions include (a) MF59 (WO 90/14837), containing 5% Squalene, 0.5% Tween 80, and 0.5% Span 85 (optionally containing various amounts of MTP-PE) formulated into submicron
- 10 particles using a microfluidizer such as Model 110Y microfluidizer (Microfluidics, Newton MA), (b) SAF, containing 10% Squalene, 0.4% Tween 80, 5% pluronic-blocked polymer L121, and thr-MDP, either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion, and (c) Ribi™ adjuvant system (RAS), (Ribi Immunochem, Hamilton, MT) containing 2% squalene, 0.2% Tween 80, and one or more
- 15 bacterial cell wall components from the group consisting of monophosphoryl lipid A, trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL + CWS (Detox™). Another class of preferred adjuvants is saponin adjuvants, such as Stimulon™ (QS-21; Aquila, Framingham, MA) or particles generated therefrom such as ISCOMs (immunostimulating complexes) and ISCOMATRIX. Other adjuvants include
- 20 Incomplete Freund's Adjuvant (IFA), cytokines, such as interleukins (IL-1, IL-2, and IL-12), macrophage colony stimulating factor (M-CSF), and tumor necrosis factor (TNF). Such adjuvants are generally available from commercial sources.

- An adjuvant can be administered with an immunogen as a single composition, or can be administered before, concurrent with or after administration of the immunogen.
- 25 Immunogen and adjuvant can be packaged and supplied in the same vial or can be packaged in separate vials and mixed before use. Immunogen and adjuvant are typically packaged with a label indicating the intended therapeutic application. If immunogen and adjuvant are packaged separately, the packaging typically includes instructions for mixing before use. The choice of an adjuvant and/or carrier depends on such factors as the
- 30 stability of the formulation containing the adjuvant, the route of administration, the dosing schedule, and the efficacy of the adjuvant for the species being vaccinated. In humans, a preferred pharmaceutically acceptable adjuvant is one that has been approved for human

such

10

20

30

particularly

can

liquid carriers, particularly for injectable solutions. Antibodies can be administered in the form of a depot injection or implant preparation which can be formulated in such a manner as to permit a sustained release of the active ingredient. An exemplary composition comprises monoclonal antibody at 5 mg/mL, formulated in aqueous buffer consisting of

5 50 mM L-histidine, 150 mM NaCl, adjusted to pH 6.0 with HCl.

Typically, compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection can also be prepared. The preparation also can be emulsified or encapsulated in liposomes or micro particles such as polylactide, polyglycolide, or

10 copolymer for enhanced adjuvant effect, as discussed above (see Langer, *Science* 249, 1527 (1990) and Hanes, *Advanced Drug Delivery Reviews* 28, 97-119 (1997)). The agents of this invention can be administered in the form of a depot injection or implant preparation which can be formulated in such a manner as to permit a sustained or pulsatile release of the active ingredient.

15 Additional formulations suitable for other modes of administration include oral, intranasal, and pulmonary formulations, suppositories, and transdermal applications.

For suppositories, binders and carriers include, for example, polyalkylene glycols or triglycerides; such suppositories can be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1%-2%. Oral formulations include

20 excipients, such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, and magnesium carbonate. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10%-95% of active ingredient, preferably 25%-70%.

Topical application can result in transdermal or intradermal delivery. Topical

25 administration can be facilitated by co-administration of the agent with cholera toxin or detoxified derivatives or subunits thereof or other similar bacterial toxins (See Glenn *et al.*, *Nature* 391, 851 (1998)). Co-administration can be achieved by using the components as a mixture or as linked molecules obtained by chemical crosslinking or expression as a fusion protein.

30 Alternatively, transdermal delivery can be achieved using a skin patch or using transferosomes (Paul *et al.*, *Eur. J. Immunol.* 25, 3521-24 (1995); Cevc *et al.*, *Biochem. Biophys. Acta* 1368, 201-15 (1998)).

# Diagnosis

## F. Methods of ~~D~~-agnosis

The invention provides methods of detecting an immune response against A $\beta$  peptide in a patient suffering from or susceptible to Alzheimer's disease. The methods are particularly useful for monitoring a course of treatment being administered to a patient.

- 5 The methods can be used to monitor both therapeutic treatment on symptomatic patients and prophylactic treatment on asymptomatic patients. The methods are useful for monitoring both active immunization (e.g., antibody produced in response to administration of immunogen) and passive immunization (e.g., measuring level of administered antibody).

10

### 1. Active Immunization

- Some methods entail determining a baseline value of an immune response in a patient before administering a dosage of agent, and comparing this with a value for the immune response after treatment. A significant increase (i.e., greater than the typical
- 15 margin of experimental error in repeat measurements of the same sample, expressed as one standard deviation from the mean of such measurements) in value of the immune response signals a positive treatment outcome (i.e., that administration of the agent has achieved or augmented an immune response). If the value for immune response does not change significantly, or decreases, a negative treatment outcome is indicated. In general, patients
- 20 undergoing an initial course of treatment with an immunogenic agent are expected to show an increase in immune response with successive dosages, which eventually reaches a plateau. Administration of agent is generally continued while the immune response is increasing. Attainment of the plateau is an indicator that the administered of treatment can be discontinued or reduced in dosage or frequency.

- 25 In other methods, a control value (i.e., a mean and standard deviation) of immune response is determined for a control population. Typically the individuals in the control population have not received prior treatment. Measured values of immune response in a patient after administering a therapeutic agent are then compared with the control value. A significant increase relative to the control value (e.g., greater than one standard deviation
- 30 from the mean) signals a positive treatment outcome. A lack of significant increase or a decrease signals a negative treatment outcome. Administration of agent is generally continued while the immune response is increasing relative to the control value. As

a plateau

indicator

before, attainment of ~~plateau~~ relative to control values in an ~~indicator~~ that the administration of treatment can be discontinued or reduced in dosage or frequency.

In other methods, a control value of immune response (e.g., a mean and standard deviation) is determined from a control population of individuals who have undergone treatment with a therapeutic agent and whose immune responses have plateaued in response to treatment. Measured values of immune response in a patient are compared with the control value. If the measured level in a patient is not significantly different (e.g., more than one standard deviation) from the control value, treatment can be discontinued. If the level in a patient is significantly below the control value, continued administration of agent is warranted. If the level in the patient persists below the control value, then a change in treatment regime, for example, use of a different adjuvant may be indicated.

In other methods, a patient who is not presently receiving treatment but has undergone a previous course of treatment is monitored for immune response to determine whether a resumption of treatment is required. The measured value of immune response in the patient can be compared with a value of immune response previously achieved in the patient after a previous course of treatment. A significant decrease relative to the previous measurement (i.e., greater than a typical margin of error in repeat measurements of the same sample) is an indication that treatment can be resumed. Alternatively, the value measured in a patient can be compared with a control value (mean plus standard deviation) determined in a population of patients after undergoing a course of treatment. Alternatively, the measured value in a patient can be compared with a control value in populations of prophylactically treated patients who remain free of symptoms of disease, or populations of therapeutically treated patients who show amelioration of disease characteristics. In all of these cases, a significant decrease relative to the control level (i.e., more than a standard deviation) is an indicator that treatment should be resumed in a patient.

The tissue sample for analysis is typically blood, plasma, serum, mucous or cerebrospinal fluid from the patient. The sample is analyzed for indication of an immune response to the amyloid component of interest, such as any form of A $\beta$  peptide. The immune response can be determined from the presence of, e.g., antibodies or T-cells that specifically bind to the component of interest, such as A $\beta$  peptide. ELISA methods of detecting antibodies specific to A $\beta$  are described in the Examples section and can be



peptide

reactive T cells

applied to other ~~peptide~~<sup>peptide</sup> antigens. Methods of detecting ~~reactive T cells~~<sup>reactive T cells</sup> are well known in the art.

## 2. Passive Immunization

5 In general, the procedures for monitoring passive immunization are similar to those for monitoring active immunization described above. However, the antibody profile following passive immunization typically shows an immediate peak in antibody concentration followed by an exponential decay. Without a further dosage, the decay approaches pretreatment levels within a period of days to months depending on the half-  
10 life of the antibody administered. For example the half-life of some human antibodies is of the order of 20 days.

In some methods, a baseline measurement of antibody to A $\beta$  in the patient is made before administration, a second measurement is made soon thereafter to determine the peak antibody level, and one or more further measurements are made at intervals to  
15 monitor decay of antibody levels. When the level of antibody has declined to baseline or a predetermined percentage of the peak less baseline (e.g., 50%, 25% or 10%), administration of a further dosage of antibody is administered. In some methods, peak or subsequent measured levels less background are compared with reference levels previously determined to constitute a beneficial prophylactic or therapeutic treatment  
20 regime in other patients. If the measured antibody level is significantly less than a reference level (e.g., less than the mean minus one standard deviation of the reference value in population of patients benefiting from treatment) administration of an additional dosage of antibody is indicated.

## 25 3. Diagnostic Kits

The invention further provides diagnostic kits for performing the diagnostic methods described above. Typically, such kits contain an agent that specifically binds to antibodies to an amyloid plaque component, such as A $\beta$ , or reacts with T-cells specific for the component. The kit can also include a label. For detection of antibodies to A $\beta$ , the  
30 label is typically in the form of labelled anti-idiotypic antibodies. For detection of antibodies, the agent can be supplied prebound to a solid phase, such as to the wells of a microtiter dish. For detection of reactive T-cells, the label can be supplied as 3H-

a typically

thymidine to measure <sup>a</sup> proliferative response. Kits also ~~typically~~ contain labelling providing directions for use of the kit. The labelling may also include a chart or other correspondence regime correlating levels of measured label with levels of antibodies to A $\beta$  or T-cells reactive with A $\beta$ . The term labelling refers to any written or recorded material

5 that is attached to, or otherwise accompanies a kit at any time during its manufacture, transport, sale or use. For example, the term labelling encompasses advertising leaflets and brochures, packaging materials, instructions, audio or video cassettes, computer discs, as well as writing imprinted directly on kits.

10

## EXAMPLES

### I. PROPHYLACTIC EFFICACY OF A $\beta$ AGAINST ALZHEIMER'S DISEASE (AD)

These examples describe administration of A $\beta$ 42 peptide to transgenic mice overexpressing APP with a mutation at position 717 (APP<sub>717V→F</sub>) that predisposes them to

15 develop Alzheimer's-like neuropathology. Production and characteristics of these mice (PDAPP mice) is described in Games et al., *Nature, supra*. These animals, in their heterozygote form, begin to deposit A $\beta$  at six months of age forward. By fifteen months of age they exhibit levels of A $\beta$  deposition equivalent to that seen in Alzheimer's disease. PDAPP mice were injected with aggregated A $\beta$ <sub>42</sub> (aggregated A $\beta$ <sub>42</sub>) or phosphate buffered

20 saline. Aggregated A $\beta$ <sub>42</sub> was chosen because of its ability to induce antibodies to multiple epitopes of A $\beta$ .

### A. METHODS

#### 1. Source of Mice

25 Thirty PDAPP heterogenic female mice were randomly divided into the following groups: 10 mice for injection with aggregated A $\beta$ 42 (one died in transit), 5 mice to be injected with PBS/adjuvant or PBS, and 10 uninjected controls. Five mice were injected with peptides dervied from the sequence of serum amyloid protein (SAP).

#### 30 2. Preparation of Immunogens

Preparation of aggregated A $\beta$ 42: two milligrams of A $\beta$ 42 (US Peptides Inc, lot K-42-12) was dissolved in 0.9 ml water and made up to 1 ml by adding 0.1 ml 10 x PBS. This was

+0

vortexed and allowed ~~to~~ incubate overnight 37° C, under which conditions the peptide aggregated. Any unused A $\beta$  was stored as a dry lyophilized powder at -20° C until the next injection.

It should be noted that when such commercially available peptides are used, the dry weights may include salt weights; weights reported in all Examples herein, unless otherwise indicated, include salt weights. Exact masses of peptide may be determined using standard assays of the preparation, such as nitrogen determination, in conjunction with the known composition.

### 3. Preparation of Injections

For each injection, 100  $\mu$ g of aggregated A $\beta$ 42 in PBS per mouse was emulsified 1:1 with Complete Freund's adjuvant (CFA) in a final volume of 400  $\mu$ l emulsion for the first immunization, followed by a boost of the same amount of immunogen in Incomplete Freund's adjuvant (IFA) at 2 weeks. Two additional doses in IFA were given at monthly intervals. The subsequent immunizations were done at monthly intervals in 500  $\mu$ l of PBS. Injections were delivered intraperitoneally (i.p.).

PBS injections followed the same schedule and mice were injected with a 1:1 mix of PBS/ Adjuvant at 400  $\mu$ l per mouse, or 500  $\mu$ l of PBS per mouse. SAP injections likewise followed the same schedule using a dose of 100  $\mu$ g per injection.

### 4. Titration of Mouse Bleeds, Tissue Preparation and Immunohistochemistry

The above methods are described *infra* in General Materials and Methods.

## B. RESULTS

PDAPP mice were injected with either aggregated A $\beta$ 42 (aggregated A $\beta$ 42), SAP peptides, or phosphate buffered saline. A group of PDAPP mice were also left as uninjected, positive controls. The titers of the mice to aggregated A $\beta$ 42 were monitored every other month from the fourth boost until the mice were one year of age. Mice were sacrificed at 13 months. At all time points examined, eight of the nine aggregated A $\beta$ 42 mice developed a high antibody titer, which remained high throughout the series of

injections (titers <sup>greater</sup> ~~greater~~ than 1/10000). The ninth mouse had a <sup>low</sup> ~~low~~, but measurable titer of approximately 1/1000 (FIG. 1, Table 3). SAPP-injected mice had titers of 1:1,000 to 1:30,000 for this immunogen with only a single mouse exceeding 1:10,000.

5

TABLE 3A  
Titers at 50% Maximal O.D.  
Aggregated A $\beta$  injected Mice

Age of PDAPP (months)	#100	#101	#102	#103	#104	#105	#106	#107	#108
4	70000	150000	15000	120000	1000	15000	50000	60000	100000
6	15000	65000	30000	55000	300	15000	15000	50000	60000
8	20000	55000	50000	50000	400	15000	18000	50000	60000
10	40000	20000	60000	50000	900	15000	50000	20000	40000
12	25000	30000	60000	40000	2700	20000	70000	25000	20000

10

TABLE 3B  
Titers at 50% Maximal O.D.  
PBS injected Mice on both Immunogens at 1/00

Age of PDAPP (months)	#113	#114	#115	#116	#117
6	<4 x bkg	<4 x bkg	<4 x bkg	<4 x bkg	<4 x bkg
10	5 x bkg	<4 x bkg	<4 x bkg	<4 x bkg	<4 x bkg
12	<4 x bkg	<4 x bkg	<4 x bkg	<4 x bkg	<4 x bkg

15

Sera from PBS-treated mice were titrated against aggregated A $\beta$ 42 at six, ten and twelve months. At a 1/100 dilution the PBS mice, when titrated against aggregated A $\beta$ 42, only exceeded 4 times background at one data point, otherwise, they were less than 4 times background at all time points (Table 3). The SAP-specific response was negligible at these time points with all titers less than 300.

20

Seven out of the nine mice in the aggregated A $\beta$ 1-42 treated group had no detectable amyloid in their brains. In contrast, brain tissue from mice in the SAP and PBS groups contained numerous amyloid deposits in the hippocampus, as well as in the frontal and cingulate cortices. The pattern of deposition was similar to that of untreated controls, with characteristic involvement of vulnerable subregions, such as the outer molecular layer of the hippocampal dentate gyrus. One mouse from the A $\beta$  1-42-injected group had a

25

amyloid

greatly reduced amyloid burden, confined to the hippocampus. An isolated plaque was identified in another A $\beta$  1-42-treated mouse.

Quantitative image analyses of the amyloid burden in the hippocampus verified the dramatic reduction achieved in the A $\beta$ 42(AN1792)-treated animals (FIG. 2). The median values of the amyloid burden for the PBS group (2.22%), and for the untreated control group (2.65%) were significantly greater than for those immunized with AN1792 (0.00%,  $p=0.0005$ ). In contrast, the median value for the group immunized with SAP peptides (SAPP) was 5.74%. Brain tissue from the untreated, control mice contained numerous A $\beta$  amyloid deposits visualized with the A $\beta$ -specific monoclonal antibody (mAb) 3D6 in the hippocampus, as well as in the retrosplenial cortex. A similar pattern of amyloid deposition was also seen in mice immunized with SAPP or PBS (FIG. 2). In addition, in these latter three groups there was a characteristic involvement of vulnerable subregions of the brain classically seen in AD, such as the outer molecular layer of the hippocampal dentate gyrus, in all three of these groups.

The brains that contained no A $\beta$  deposits were also devoid of neuritic plaques that are typically visualized in PDAPP mice with the human APP antibody 8E5. All of brains from the remaining groups (SAP-injected, PBS and uninjected mice) had numerous neuritic plaques typical of untreated PDAPP mice. A small number of neuritic plaques were present in one mouse treated with AN1792, and a single cluster of dystrophic neurites was found in a second mouse treated with AN1792. Image analyses of the hippocampus, and shown in FIG. 3, demonstrated the virtual elimination of dystrophic neurites in AN1792-treated mice (median 0.00%) compared to the PBS recipients (median 0.28%,  $p = 0.0005$ ).

Astrocytosis characteristic of plaque-associated inflammation was also absent in the brains of the A $\beta$ 1-42 injected group. The brains from the mice in the other groups contained abundant and clustered GFAP-positive astrocytes typical of A $\beta$  plaque-associated gliosis. A subset of the GFAP-reacted slides were counter-stained with Thioflavin S to localize the A $\beta$  deposits. The GFAP-positive astrocytes were associated with A $\beta$  plaques in the SAP, PBS and untreated controls. No such association was found in the plaque-negative A $\beta$ 1-42 treated mice, while minimal plaque-associated gliosis was identified in one mouse treated with AN1792.

analyses,

Image analysis shown in FIG. 4 for the retrosplenial cortex, verified that the reduction in astrocytosis was significant with a median value of 1.56% for those treated with AN1792 versus median values greater than 6% for groups immunized with SAP peptides, PBS or untreated ( $p=0.0017$ )

- 5 Evidence from a subset of the A $\beta$ 1-42- and PBS-injected mice indicated plaque-associated MHC II immunoreactivity was absent in the A $\beta$ 1-42 injected mice, consistent with lack of an A $\beta$ -related inflammatory response.

- Sections of the mouse brains were also reacted with a mAb specific with a monoclonal antibody specific for MAC-1, a cell surface protein. MAC-1 (CD11b) is an  
 10 integrin family member and exists as a heterodimer with CD18. The CD11b/CD18 complex is present on monocytes, macrophages, neutrophils and natural killer cells (Mak and Simard). The resident MAC-1-reactive cell type in the brain is likely to be microglia based on similar phenotypic morphology in MAC-1 immunoreacted sections. Plaque-associated MAC-1 labeling was lower in the brains of mice treated with AN1792  
 15 compared to the PBS control group, a finding consistent with the lack of an A $\beta$ -induced inflammatory response.

### C. CONCLUSION

- The lack of A $\beta$  plaques and reactive neuronal and gliotic changes in the brains of  
 20 the A $\beta$ 1-42-injected mice indicate that no or extremely little amyloid was deposited in their brains, and pathological consequences, such as gliosis and neuritic pathology, were absent. PDAPP mice treated with A $\beta$ 1-42 show essentially the same lack of pathology as control nontransgenic mice. Therefore, A $\beta$ 1-42 injections are highly effective in the prevention of deposition or clearance of human A $\beta$  from brain tissue, and elimination of  
 25 subsequent neuronal and inflammatory degenerative changes. Thus, administration of A $\beta$  peptide can have both preventative and therapeutic benefit in prevention of AD.

### II. DOSE RESPONSE STUDY

- Groups of five-week old, female Swiss Webster mice (N=6 per group) were  
 30 immunized with 300, 100, 33, 11, 3.7, 1.2, 0.4, or 0.13 ug of A $\beta$  formulated in CFA/IFA administered intraperitoneally. Three doses were given at biweekly intervals followed by a fourth dose one month later. The first dose was emulsified with CFA and the remaining

emulsified with

following

09/724570

doses were ~~emulsified~~ with IFA. Animals were bled 4-7 days ~~following~~ each immunization starting after the second dose for measurement of antibody titers. Animals in a subset of three groups, those immunized with 11, 33, or 300  $\mu\text{g}$  of antigen, were additionally bled at approximately monthly intervals for four months following the fourth immunization to monitor the decay of the antibody response across a range of doses of immunogenic formulations. These animals received a final fifth immunization at seven months after study initiation. They were sacrificed one week later to measure antibody responses to AN1792 and to perform toxicological analyses.

10 A declining dose response was observed from 300 to 3.7  $\mu\text{g}$  with no response at the two lowest doses. Mean antibody titers are about 1:1000 after 3 doses and about 1:10,000 after 4 doses of 11-300  $\mu\text{g}$  of antigen (see FIG. 5).

Antibody titers rose dramatically for all but the lowest dose group following the third immunization with increases in GMTs ranging from 5- to 25-fold. Low antibody responses were then detectable for even the 0.4  $\mu\text{g}$  recipients. The 1.2 and 3.7  $\mu\text{g}$  groups had comparable titers with GMTs of about 1000 and the highest four doses clustered together with GMTs of about 25,000, with the exception of the 33  $\mu\text{g}$  dose group with a lower GMT of 3000. Following the fourth immunization, the titer increase was more modest for most groups. There was a clear dose response across the lower antigen dose groups from 0.14  $\mu\text{g}$  to 11  $\mu\text{g}$  ranging from no detectable antibody for recipients of 0.14  $\mu\text{g}$  to a GMT of 36,000 for recipients of 11  $\mu\text{g}$ . Again, titers for the four highest dose groups of 11 to 300  $\mu\text{g}$  clustered together. Thus following two immunizations, the antibody titer was dependent on the antigen dose across the broad range from 0.4 to 300  $\mu\text{g}$ . By the third immunization, titers of the highest four doses were all comparable and they remained at a plateau after an additional immunization.

One month following the fourth immunization, titers were 2- to 3-fold higher in the 300  $\mu\text{g}$  group than those measured from blood drawn five days following the immunization (FIG. 6). This observation suggests that the peak anamnestic antibody response occurred later than 5 days post-immunization. A more modest (50%) increase was seen at this time in the 33  $\mu\text{g}$  group. In the 300  $\mu\text{g}$  dose group at two months following the last dose, GMTs declined steeply by about 70%. After another month, the decline was less steep at 45% (100  $\mu\text{g}$ ) and about 14% for the 33 and 11  $\mu\text{g}$  doses. Thus,

*circulating*

the rate of decline in ~~circulating~~ antibody titers following cessation of immunization appears to be biphasic with a steep decline the first month following peak response followed by a more modest rate of decrease thereafter.

- 5 The antibody titers and the kinetics of the response of these Swiss Webster mice are similar to those of young heterozygous PDAPP transgenic mice immunized in a parallel manner. Dosages effective to induce an immune response in humans are typically similar to dosages effective in mice.

### III. SCREEN FOR THERAPEUTIC EFFICACY AGAINST ESTABLISHED AD

- 10 This assay is designed to test immunogenic agents for activity in arresting or reversing neuropathologic characteristics of AD in aged animals. Immunizations with 42 amino acid long A $\beta$  (AN1792) were begun at a time point when amyloid plaques are already present in the brains of the PDAPP mice.

- Over the time course used in this study, untreated PDAPP mice develop a number  
15 of neurodegenerative changes that resemble those found in AD (*Games et al., supra* and Johnson-Wood et al., *Proc. Natl. Acad. Sci. USA* 94, 1550-1555 (1997)). The deposition of A $\beta$  into amyloid plaques is associated with a degenerative neuronal response consisting of aberrant axonal and dendritic elements, called dystrophic neurites. Amyloid deposits that are surrounded by and contain dystrophic neurites called neuritic plaques. In both AD  
20 and the PDAPP mouse, dystrophic neurites have a distinctive globular structure, are immunoreactive with a panel of antibodies recognizing APP and cytoskeletal components, and display complex subcellular degenerative changes at the ultrastructural level. These characteristics allow for disease-relevant, selective and reproducible measurements of neuritic plaque formation in the PDAPP brains. The dystrophic neuronal component of  
25 PDAPP neuritic plaques is easily visualized with an antibody specific for human APP (monoclonal antibody 8E5), and is readily measurable by computer-assisted image analysis. Therefore, in addition to measuring the effects of AN1792 on amyloid plaque formation, we monitored the effects of this treatment on the development of neuritic dystrophy.

- 30 Astrocytes and microglia are non-neuronal cells that respond to and reflect the degree of neuronal injury. GFAP-positive astrocytes and MHC II-positive microglia are commonly observed in AD, and their activation increases with the severity of the disease.



monitored

astrocytosis

09/724570

Therefore, we also ~~monitored~~ the development of reactive ~~astrocytosis~~ and microgliosis in the AN1792-treated mice.

#### A. Materials and Methods

5 Forty-eight, heterozygous female PDAPP mice, 11 to 11.5 months of age, obtained from Charles River, were randomly divided into two groups: 24 mice to be immunized with 100  $\mu$ g of AN1792 and 24 mice to be immunized with PBS, each combined with Freund's adjuvant. The AN1792 and PBS groups were again divided when they reached ~15 months of age. At 15 months of age approximately half of each group of the  
10 AN1792- and PBS-treated animals were euthanized (n=10 and 9, respectively), the remainder continued to receive immunizations until termination at ~18 months (n=9 and 12, respectively). A total of 8 animals (5 AN1792, 3 PBS) died during the study. In addition to the immunized animals, one-year old (n=10), 15-month old (n=10) and 18-month old (n=10) untreated PDAPP mice were included for comparison in the ELISAs to  
15 measure A $\beta$  and APP levels in the brain; the one-year old animals were also included in the immunohistochemical analyses.

Methodology was as in Example I, unless otherwise indicated. US Peptides lot 12 and California Peptides lot ME0339 of AN1792 were used to prepare the antigen for the six immunizations administered prior to the 15-month time point. California Peptides lots  
20 ME0339 and ME0439 were used for the three additional immunizations administered between 15 and 18 months.

For immunizations, 100  $\mu$ g of AN1792 in 200  $\mu$ l PBS or PBS alone was emulsified 1:1 (vol:vol) with Complete Freund's adjuvant (CFA) or Incomplete Freund's adjuvant (IFA) or PBS in a final volume of 400  $\mu$ l. The first immunization was delivered with CFA  
25 as adjuvant, the next four doses were given with IFA and the final four doses with PBS alone without added adjuvant. A total of nine immunizations were given over the seven-month period on a two-week schedule for the first three doses followed by a four-week interval for the remaining injections. The four-month treatment group, euthanized at 15 months of age, received only the first 6 immunizations.

30

#### B. Results

##### 1. Effects of A $\beta$ 42 (AN1792) Treatment on Amyloid Burden

results of

amyloid

The ~~result of~~ AN1792 treatment on cortical amyloid burden determined by quantitative image analysis are shown in FIG. 7. The median value of cortical amyloid burden was 0.28% in a group of untreated 12-month old PDAPP mice, a value representative of the plaque load in mice at the study's initiation. At 18 months, the amyloid burden increased over 17-fold to 4.87% in PBS-treated mice, while AN1792-treated mice had a greatly reduced amyloid burden of only 0.01%, notably less than the 12-month untreated and both the 15- and 18-month PBS-treated groups. The amyloid burden was significantly reduced in the AN1792 recipients at both 15 (96% reduction;  $p=0.003$ ) and 18 (>99% reduction;  $p=0.0002$ ) months.

Typically, cortical amyloid deposition in PDAPP mice initiates in the frontal and retrosplenial cortices (RSC) and progresses in a ventral-lateral direction to involve the temporal and entorhinal cortices (EC). Little or no amyloid was found in the EC of 12 month-old mice, the approximate age at which AN1792 was first administered. After 4 months of AN1792 treatment, amyloid deposition was greatly diminished in the RSC, and the progressive involvement of the EC was entirely eliminated by AN1792 treatment. The latter observation showed that AN1792 completely halted the progression of amyloid that would normally invade the temporal and ventral cortices, as well as arrested or possibly reversed deposition in the RSC.

The profound effects of AN1792 treatment on developing cortical amyloid burden in the PDAPP mice are further demonstrated by the 18-month group, which had been treated for seven months. A near complete absence of cortical amyloid was found in the AN1792-treated mouse, with a total lack of diffuse plaques, as well as a reduction in compacted deposits.

## 2. A $\beta$ 42 (AN1792) Treatment-associated Cellular and Morphological Changes

A population of A $\beta$ -positive cells was found in brain regions that typically contain amyloid deposits. Remarkably, in several brains from AN1792 recipients, very few or no extracellular cortical amyloid plaques were found. Most of the A $\beta$  immunoreactivity appeared to be contained within cells with large lobular or clumped soma. Phenotypically, these cells resembled activated microglia or monocytes. They were immunoreactive with antibodies recognizing ligands expressed by activated monocytes and microglia (MHC II and CD11b) and were occasionally associated with the wall or

vessels.

lumen of blood ~~vessels~~. Comparison of near-adjacent sections labeled with A $\beta$  and MHC II-specific antibodies revealed that similar patterns of these cells were recognized by both classes of antibodies. Detailed examination of the AN1792-treated brains revealed that the MHC II-positive cells were restricted to the vicinity of the limited amyloid remaining in these animals. Under the fixation conditions employed, the cells were not immunoreactive with antibodies that recognize T cell (CD3, CD3e) or B cell (CD45RA, CD45RB) ligands or leukocyte common antigen (CD45), but were reactive with an antibody recognizing leukosialin (CD43) which cross-reacts with monocytes. No such cells were found in any of the PBS-treated mice.

PDAPP mice invariably develop heavy amyloid deposition in the outer molecular layer of the hippocampal dentate gyrus. The deposition forms a distinct streak within the perforant pathway, a subregion that classically contains amyloid plaques in AD. The characteristic appearance of these deposits in PBS-treated mice resembled that previously characterized in untreated PDAPP mice. The amyloid deposition consisted of both diffuse and compacted plaques in a continuous band. In contrast, in a number of brains from AN1792-treated mice this pattern was drastically altered. The hippocampal amyloid deposition no longer contained diffuse amyloid, and the banded pattern was completely disrupted. Instead, a number of unusual punctate structures were present that are reactive with anti-A $\beta$  antibodies, several of which appeared to be amyloid-containing cells.

MHC II-positive cells were frequently observed in the vicinity of extracellular amyloid in AN1792-treated animals. The pattern of association of A $\beta$ -positive cells with amyloid was very similar in several brains from AN1792-treated mice. The distribution of these monocytic cells was restricted to the proximity of the deposited amyloid and was entirely absent from other brain regions devoid of A $\beta$  plaques.

Quantitative image analysis of MHC II and MAC I-labeled sections revealed a trend towards increased immunoreactivity in the RSC and hippocampus of AN1792-treated mice compared to the PBS group which reached significance with the measure of MAC 1 reactivity in hippocampus.

These results are indicative of active, cell-mediated removal of amyloid in plaque-bearing brain regions.

## AN1792

3. ~~AN1792~~ Effects on A $\beta$  Levels: ELISA Determinations

## (a) Cortical Levels

In untreated PDAPP mice, the median level of total A $\beta$  in the cortex at 12 months was 1,600 ng/g, which increased to 8,700 ng/g by 15 months (Table 4). At 18 months the value was 22,000 ng/g, an increase of over 10-fold during the time course of the experiment. PBS-treated animals had 8,600 ng/g total A $\beta$  at 15 months which increased to 19,000 ng/g at 18 months. In contrast, AN1792-treated animals had 81% less total A $\beta$  at 15 months (1,600 ng/g) than the PBS-immunized group. Significantly less ( $p=0.0001$ ) total A $\beta$  (5,200 ng/g) was found at 18 months when the AN1792 and PBS groups were compared (Table 4), representing a 72% reduction in the A $\beta$  that would otherwise be present. Similar results were obtained when cortical levels of A $\beta$ 42 were compared, namely that the AN1792-treated group contained much less A $\beta$ 42, but in this case the differences between the AN1792 and PBS groups were significant at both 15 months ( $p=0.04$ ) and 18 months ( $p=0.0001$ , Table 4).

Table 4: Median A $\beta$  Levels (ng/g) in Cortex

Age	UNTREATED			PBS			AN1792		
	Total	A $\beta$ 42	(n)	Total	A $\beta$ 42	(n)	Total	A $\beta$ 42	(n)
12	1,600	1,300	(10)						
15	8,700	8,300	(10)	8,600	7,200	(9)	1,600	1,300*	(10)
18	22,200	18,500	(10)	19,000	15,900	(12)	5,200**	4,000**	(9)

\*  $p = 0.0412$ \*\*  $p = 0.0001$ 

## (b) Hippocampal Levels

In untreated PDAPP mice, median hippocampal levels of total A $\beta$  at twelve months of age were 15,000 ng/g which increased to 51,000 ng/g at 15 months and further to 81,000 ng/g at 18 months (Table 5). Similarly, PBS immunized mice showed values of 40,000 ng/g and 65,000 ng/g at 15 months and 18 months, respectively. AN1792 immunized animals exhibited less total A $\beta$ , specifically 25,000 ng/g and 51,000 ng/g at the respective 15-month and 18-month timepoints. The 18-month AN1792-treated group value was significantly lower than that of the PBS treated group ( $p=0.0105$ ; Table 5). Measurement of A $\beta$ 42 gave the same pattern of results, namely that levels in the AN1792-treated group were significantly lower than in the PBS group (39,000 ng/g vs. 57,000 ng/g, respectively;  $p=0.002$ ) at the 18-month evaluation (Table 3).

values

normally

In all cases the APP ~~values~~ remained within the ranges that are normally found in PDAPP mice.

## 5 5. Effects of AN1792 Treatment on Neurodegenerative and Gliotic Pathology

Neuritic plaque burden was significantly reduced in the frontal cortex of AN1792-treated mice compared to the PBS group at both 15 (84%;  $p=0.03$ ) and 18 (55%;  $p=0.01$ ) months of age (FIG. 8). The median value of the neuritic plaque burden increased from 0.32% to 0.49% in the PBS group between 15 and 18 months of age. This contrasted with the greatly reduced development of neuritic plaques in the AN1792 group, with median neuritic plaque burden values of 0.05% and 0.22%, in the 15 and 18 month groups, respectively.

Immunizations with AN1792 seemed well tolerated and reactive astrocytosis was also significantly reduced in the RSC of AN1792-treated mice when compared to the PBS group at both 15 (56%;  $p=0.011$ ) and 18 (39%;  $p=0.028$ ) months of age (FIG. 9). Median values of the percent of astrocytosis in the PBS group increased between 15 and 18 months from 4.26% to 5.21%. AN1792-treatment suppressed the development of astrocytosis at both time points to 1.89% and 3.2%, respectively. This suggests the neuropil was not being damaged by the clearance process.

20

## 6. Antibody Responses

As described above, eleven-month old, heterozygous PDAPP mice ( $N=24$ ) received a series of 5 immunizations of 100  $\mu$ g of AN1792 emulsified with Freund's adjuvant and administered intraperitoneally at weeks 0, 2, 4, 8, and 12, and a sixth immunization with PBS alone (no Freund's adjuvant) at week 16. As a negative control, a parallel set of 24 age-matched transgenic mice received immunizations of PBS emulsified with the same adjuvants and delivered on the same schedule. Animals were bled within three to seven days following each immunization starting after the second dose. Antibody responses to AN1792 were measured by ELISA. Geometric mean titers (GMT) for the animals that were immunized with AN1792 were approximately 1,900, 7,600, and 45,000 following the second, third and last (sixth) doses respectively. No A $\beta$ -specific antibody was measured in control animals following the sixth immunization.

Approximately

for

Approximately one-half of the animals were treated for an additional three months, receiving immunizations at about 20, 24 and 27 weeks. Each of these doses was delivered in PBS vehicle alone without Freund's adjuvant. Mean antibody titers remained unchanged over this time period. In fact, antibody titers appeared to remain stable from the fourth to the eighth bleed corresponding to a period covering the fifth to the ninth injections.

To determine if the A $\beta$ -specific antibodies elicited by immunization that were detected in the sera of AN1792-treated mice were also associated with deposited brain amyloid, a subset of sections from the AN1792- and PBS-treated mice were reacted with an antibody specific for mouse IgG. In contrast to the PBS group, A $\beta$  plaques in AN1792-treated brains were coated with endogenous IgG. This difference between the two groups was seen in both 15- and 18-month groups. Particularly striking was the lack of labeling in the PBS group, despite the presence of a heavy amyloid burden in these mice. These results show that immunization with a synthetic A $\beta$  protein generates antibodies that recognize and bind in vivo to the A $\beta$  in amyloid plaques.

#### 7. Cellular-Mediated Immune Responses

Spleens were removed from nine AN1792-immunized and 12 PBS-immunized 18-month old PDAPP mice 7 days after the ninth immunization. Splenocytes were isolated and cultured for 72 h in the presence of A $\beta$ 40, A $\beta$ 42, or A $\beta$ 40-1 (reverse order protein). The mitogen Con A served as a positive control. Optimum responses were obtained with >1.7  $\mu$ M protein. Cells from all nine AN1792-treated animals proliferated in response to either A $\beta$ 1-40 or A $\beta$ 1-42 protein, with equal levels of incorporation for both proteins (FIG. 10, Upper Panel). There was no response to the A $\beta$ 40-1 reverse protein. Cells from control animals did not respond to any of the A $\beta$  proteins (FIG. 10, Lower Panel).

#### C. Conclusion

The results of this study show that AN1792 immunization of PDAPP mice possessing existing amyloid deposits slows and prevents progressive amyloid deposition and retard consequential neuropathologic changes in the aged PDAPP mouse brain. Immunizations with AN1792 essentially halted amyloid developing in structures that

succumb

administration of

09/724570

would normally succumb to amyloidosis. Thus, administration of A $\beta$  peptide has therapeutic benefit in the treatment of AD.

#### IV. SCREEN OF A $\beta$ FRAGMENTS

5 100 PDAPP mice age 9-11 months were immunized with 9 different regions of APP and A $\beta$  to determine which epitopes convey the efficacious response. The 9 different immunogens and one control are injected i.p. as described above. The immunogens include four human A $\beta$  peptide conjugates 1-12, 13-28, 32-42, 1-5, all coupled to sheep anti-mouse IgG via a cystine link; an APP polypeptide amino acids 592-695, aggregated  
10 human A $\beta$  1-40, and aggregated human A $\beta$  25-35, and aggregated rodent A $\beta$ 42. Aggregated A $\beta$ 42 and PBS were used as positive and negative controls, respectively. Ten mice were used per treatment group. Titers were monitored as above and mice were euthanized at the end of 4 months of injections. Histochemistry, A $\beta$  levels, and toxicology analysis was determined post mortem.

15

##### A. Materials and Methods

##### 1. Preparation of Immunogens

Preparation of coupled A $\beta$  peptides: four human A $\beta$  peptide conjugates (amino acid residues 1-5, 1-12, 13-28, and 33-42, each conjugated to sheep anti-mouse IgG) were  
20 prepared by coupling through an artificial cysteine added to the A $\beta$  peptide using the crosslinking reagent sulfo-EMCS. The A $\beta$  peptide derivatives were synthesized with the following final amino acid sequences. In each case, the location of the inserted cysteine residue is indicated by underlining. The A $\beta$ 13-28 peptide derivative also had two glycine residues added prior to the carboxyl terminal cysteine as indicated.

25

A $\beta$ 1-12 peptide	NH <sub>2</sub> -DAEFRHDSGYEVC-COOH (SEQ ID NO: 30)
A $\beta$ 1-5 peptide	NH <sub>2</sub> -DAEFR <u>C</u> -COOH (SEQ ID NO: 31)
A $\beta$ 33-42 peptide	NH <sub>2</sub> - <u>C</u> -amino-heptanoic acid-GLMVGGVVIA-COOH (SEQ ID NO: 32)
30 A $\beta$ 13-28 peptide	Ac-NH-HHQLVFFAEDVGSNKG <u>G</u> C-COOH (SEQ ID NO: 33)

prepare for

To ~~prepare for~~ the coupling reaction, ten mg of sheep anti-mouse IgG (Jackson ImmunoResearch Laboratories) was dialyzed overnight against 10 mM sodium borate buffer, pH 8.5. The dialyzed antibody was then concentrated to a volume of 2 mL using an Amicon Centriprep tube. Ten mg sulfo-EMCS

- 5 [N ( $\gamma$ -maleimidocaproyloxy) succinimide] (Molecular Sciences Co.) was dissolved in one mL deionized water. A 40-fold molar excess of sulfo-EMCS was added dropwise with stirring to the sheep anti-mouse IgG and then the solution was stirred for an additional ten min. The activated sheep anti-mouse IgG was purified and buffer
- 10 exchanged by passage over a 10 mL gel filtration column (Pierce Presto Column, obtained from Pierce Chemicals) equilibrated with 0.1 M NaPO<sub>4</sub>, 5 mM EDTA, pH 6.5. Antibody containing fractions, identified by absorbance at 280 nm, were pooled and diluted to a concentration of approximately 1 mg/mL, using 1.4 mg per OD as the extinction
- 15 coefficient. A 40-fold molar excess of A $\beta$  peptide was dissolved in 20 mL of 10 mM NaPO<sub>4</sub>, pH 8.0, with the exception of the A $\beta$ 33-42 peptide for which 10 mg was first dissolved in 0.5 mL of DMSO and then diluted to 20 mL with the 10 mM NaPO<sub>4</sub> buffer.
- The peptide solutions were each added to 10 mL of activated sheep anti-mouse IgG and rocked at room temperature for 4 hr. The resulting conjugates were concentrated to a final volume of less than 10 mL using an Amicon Centriprep tube and then dialyzed against
- 20 PBS to buffer exchange the buffer and remove free peptide. The conjugates were passed through 0.22  $\mu$ m-pore size filters for sterilization and then aliquoted into fractions of 1 mg and stored frozen at -20°C. The concentrations of the conjugates were determined using the BCA protein assay (Pierce Chemicals) with horse IgG for the standard curve.
- Conjugation was documented by the molecular weight increase of the conjugated peptides relative to that of the activated sheep anti-mouse IgG. The A $\beta$  1-5 sheep anti-mouse
- 25 conjugate was a pool of two conjugations, the rest were from a single preparation.

## 2. Preparation of aggregated A $\beta$ peptides

- Human 1-40 (AN1528; California Peptides Inc., Lot ME0541), human 1-42 (AN1792; California Peptides Inc., Lots ME0339 and ME0439), human 25-35, and rodent
- 30 1-42 (California Peptides Inc., Lot ME0218) peptides were freshly solubilized for the preparation of each set of injections from lyophilized powders that had been stored desiccated at -20°C. For this purpose, two mg of peptide were added to 0.9 ml of



the  
 deionized water and ~~the~~ mixture was vortexed to generate a relatively uniform solution or suspension. Of the four, AN1528 was the only peptide soluble at this step. A 100  $\mu$ l aliquot of 10X PBS (1X PBS: 0.15 M NaCl, 0.01 M sodium phosphate, pH 7.5) was then added at which point AN1528 began to precipitate. The suspension was vortexed again  
 5 and incubated overnight at 37°C for use the next day.

Preparation of the pBx6 protein: An expression plasmid encoding pBx6, a fusion protein consisting of the 100-amino acid bacteriophage MS-2 polymerase N-terminal leader sequence followed by amino acids 592-695 of APP ( $\beta$ APP) was constructed as described by Oltersdorf et al., J. Biol. Chem. 265, 4492-4497 (1990). The  
 10 plasmid was transfected into E. coli and the protein was expressed after induction of the promoter. The bacteria were lysed in 8M urea and pBx6 was partially purified by preparative SDS PAGE. Fractions containing pBx6 were identified by Western blot using a rabbit anti-pBx6 polyclonal antibody, pooled, concentrated using an Amicon Centriprep tube and dialysed against PBS. The purity of the preparation, estimated by Coomassie  
 15 Blue stained SDS PAGE, was approximately 5 to 10%.

## B. Results and Discussion

### 1. Study Design

One hundred male and female, nine- to eleven-month old heterozygous PDAPP  
 20 transgenic mice were obtained from Charles River Laboratory and Taconic Laboratory. The mice were sorted into ten groups to be immunized with different regions of A $\beta$  or APP combined with Freund's adjuvant. Animals were distributed to match the gender, age, parentage and source of the animals within the groups as closely as possible. The immunogens included four A $\beta$  peptides derived from the human sequence, 1-5, 1-12, 13-  
 25 28, and 33-42, each conjugated to sheep anti-mouse IgG; four aggregated A $\beta$  peptides, human 1-40 (AN1528), human 1-42 (AN1792), human 25-35, and rodent 1-42; and a fusion polypeptide, designated as pBx6, containing APP amino acid residues 592-695. A tenth group was immunized with PBS combined with adjuvant as a control.

For each immunization, 100  $\mu$ g of each A $\beta$  peptide in 200  $\mu$ l PBS or 200  $\mu$ g of  
 30 the APP derivative pBx6 in the same volume of PBS or PBS alone was emulsified 1:1 (vol:vol) with Complete Freund's adjuvant (CFA) in a final volume of 400  $\mu$ l for the first immunization, followed by a boost of the same amount of immunogen in Incomplete

(IFA)

PBS

09/724570

Freund's adjuvant (H<sub>2</sub>O) for the subsequent four doses and with PBS for the final dose. Immunizations were delivered intraperitoneally on a biweekly schedule for the first three doses, then on a monthly schedule thereafter. Animals were bled four to seven days following each immunization starting after the second dose for the measurement of antibody titers. Animals were euthanized approximately one week after the final dose.

## 2. A $\beta$ and APP Levels in the Brain

Following about four months of immunization with the various A $\beta$  peptides or the APP derivative, brains were removed from saline-perfused animals. One hemisphere was prepared for immunohistochemical analysis and the second was used for the quantitation of A $\beta$  and APP levels. To measure the concentrations of various forms of beta amyloid peptide and amyloid precursor protein, the hemisphere was dissected and homogenates of the hippocampal, cortical, and cerebellar regions were prepared in 5 M guanidine. These were diluted and the level of amyloid or APP was quantitated by comparison to a series of dilutions of standards of A $\beta$  peptide or APP of known concentrations in an ELISA format.

The median concentration of total A $\beta$  for the control group immunized with PBS was 5.8-fold higher in the hippocampus than in the cortex (median of 24,318 ng/g hippocampal tissue compared to 4,221 ng/g for the cortex). The median level in the cerebellum of the control group (23.4 ng/g tissue) was about 1,000-fold lower than in the hippocampus. These levels are similar to those that we have previously reported for heterozygous PDAPP transgenic mice of this age (Johnson-Woods et al., 1997, supra).

For the cortex, a subset of treatment groups had median total A $\beta$  and A $\beta$ 1-42 levels which differed significantly from those of the control group ( $p < 0.05$ ), those animals receiving AN1792, rodent A $\beta$ 1-42 or the A $\beta$ 1-5 peptide conjugate as shown in FIG. 11. The median levels of total A $\beta$  were reduced by 75%, 79% and 61%, respectively, compared to the control for these treatment groups. There were no discernable correlations between A $\beta$ -specific antibody titers and A $\beta$  levels in the cortical region of the brain for any of the groups.

In the hippocampus, the median reduction of total A $\beta$  associated with AN1792 treatment (46%,  $p = 0.0543$ ) was not as great as that observed in the cortex (75%,  $p = 0.0021$ ). However, the magnitude of the reduction was far greater in the hippocampus

net

than in the cortex, a ~~reduction~~ reduction of 11,186 ng/g tissue in the hippocampus versus 3,171 ng/g tissue in the cortex. For groups of animals receiving rodent A $\beta$ 1-42 or A $\beta$ 1-5, the median total A $\beta$  levels were reduced by 36% and 26%, respectively. However, given the small group sizes and the high variability of the amyloid peptide levels from animal to animal within both groups, these reductions were not significant. When the levels of A $\beta$ 1-42 were measured in the hippocampus, none of the treatment-induced reductions reached significance. Thus, due to the smaller A $\beta$  burden in the cortex, changes in this region are a more sensitive indicator of treatment effects. The changes in A $\beta$  levels measured by ELISA in the cortex are similar, but not identical, to the results from the immunohistochemical analysis (see below).

Total A $\beta$  was also measured in the cerebellum, a region typically minimally affected with AD pathology. None of the median A $\beta$  concentrations of any of the groups immunized with the various A $\beta$  peptides or the APP derivative differed from that of the control group in this region of the brain. This result suggests that non-pathological levels of A $\beta$  are unaffected by treatment.

APP concentration was also determined by ELISA in the cortex and cerebellum from treated and control mice. Two different APP assays were utilized. The first, designated APP- $\alpha$ /FL, recognizes both APP-alpha ( $\alpha$ , the secreted form of APP which has been cleaved within the A $\beta$  sequence), and full-length forms (FL) of APP, while the second recognizes only APP- $\alpha$ . In contrast to the treatment-associated diminution of A $\beta$  in a subset of treatment groups, the levels of APP were unchanged in all of the treated compared to the control animals. These results indicate that the immunizations with A $\beta$  peptides are not depleting APP; rather the treatment effect is specific to A $\beta$ .

In summary, total A $\beta$  and A $\beta$ 1-42 levels were significantly reduced in the cortex by treatment with AN1792, rodent A $\beta$ 1-42 or A $\beta$ 1-5 conjugate. In the hippocampus, total A $\beta$  was significantly reduced only by AN1792 treatment. No other treatment-associated changes in A $\beta$  or APP levels in the hippocampal, cortical or cerebellar regions were significant.

## 2. Histochemical Analyses

from

Brains ~~from~~ a subset of six groups were prepared for immunohistochemical analysis, three groups immunized with the A $\beta$  peptide conjugates A $\beta$ 1-5, A $\beta$ 1-12, and A $\beta$ 13-28; two groups immunized with the full length A $\beta$  aggregates AN1792 and AN1528 and the PBS-treated control group. The results of image analyses of the amyloid burden in brain sections from these groups are shown in FIG. 12. There were significant reductions of amyloid burden in the cortical regions of three of the treatment groups versus control animals. The greatest reduction of amyloid burden was observed in the group receiving AN1792 where the mean value was reduced by 97% ( $p = 0.001$ ). Significant reductions were also observed for those animals treated with AN1528 (95%,  $p = 0.005$ ) and the A $\beta$ 1-5 peptide conjugate (67%,  $p = 0.02$ ).

The results obtained by quantitation of total A $\beta$  or A $\beta$ 1-42 by ELISA and amyloid burden by image analysis differ to some extent. Treatment with AN1528 had a significant impact on the level of cortical amyloid burden when measured by quantitative image analysis but not on the concentration of total A $\beta$  in the same region when measured by ELISA. The difference between these two results is likely to be due to the specificities of the assays. Image analysis measures only insoluble A $\beta$  aggregated into plaques. In contrast, the ELISA measures all forms of A $\beta$ , both soluble and insoluble, monomeric and aggregated. Since the disease pathology is thought to be associated with the insoluble plaque-associated form of A $\beta$ , the image analysis technique may have more sensitivity to reveal treatment effects. However since the ELISA is a more rapid and easier assay, it is very useful for screening purposes. Moreover it may reveal that the treatment-associated reduction of A $\beta$  is greater for plaque-associated than total A $\beta$ .

To determine if the A $\beta$ -specific antibodies elicited by immunization in the treated animals reacted with deposited brain amyloid, a subset of the sections from the treated animals and the control mice were reacted with an antibody specific for mouse IgG. In contrast to the PBS group, A $\beta$ -containing plaques were coated with endogenous IgG for animals immunized with the A $\beta$  peptide conjugates A $\beta$ 1-5, A $\beta$ 1-12, and A $\beta$ 13-28; and the full length A $\beta$  aggregates AN1792 and AN1528. Brains from animals immunized with the other A $\beta$  peptides or the APP peptide pBx6 were not analyzed by this assay.

### 3. Measurement of Antibody Titers

bled

Mice were ~~used~~ bled four to seven days following each immunization starting after the second immunization, for a total of five bleeds. Antibody titers were measured as A $\beta$ 1-42-binding antibody using a sandwich ELISA with plastic multi-well plates coated with A $\beta$ 1-42. As shown in FIG. 13, peak antibody titers were elicited following the fourth dose for those four immunization formulations which elicited the highest titers of AN1792-specific antibodies: AN1792 (peak GMT: 94,647), AN1528 (peak GMT: 88,231), A $\beta$ 1-12 conjugate (peak GMT: 47,216) and rodent A $\beta$ 1-42 (peak GMT: 10,766). Titers for these groups declined somewhat following the fifth and sixth doses. For the remaining five immunogens, peak titers were reached following the fifth or the sixth dose and these were of much lower magnitude than those of the four highest titer groups: A $\beta$ 1-5 conjugate (peak GMT: 2,356), pBx6 (peak GMT: 1,986), A $\beta$ 13-28 conjugate (peak GMT: 1,183), A $\beta$ 33-42 conjugate (peak GMT: 658), A $\beta$ 25-35 (peak GMT: 125). Antibody titers were also measured against the homologous peptides using the same ELISA sandwich format for a subset of the immunogens, those groups immunized with A $\beta$ 1-5, A $\beta$ 13-28, A $\beta$ 25-35, A $\beta$ 33-42 or rodent A $\beta$ 1-42. These titers were about the same as those measured against A $\beta$ 1-42 except for the rodent A $\beta$ 1-42 immunogen in which case antibody titers against the homologous immunogen were about two-fold higher. The magnitude of the AN1792-specific antibody titer of individual animals or the mean values of treatment groups did not correlate with efficacy measured as the reduction of A $\beta$  in the cortex.

#### 4. Lymphoproliferative Responses

A $\beta$ -dependent lymphoproliferation was measured using spleen cells harvested approximately one week following the final, sixth, immunization. Freshly harvested cells, 105 per well, were cultured for 5 days in the presence of A $\beta$ 1-40 at a concentration of 5  $\mu$ M for stimulation. Cells from a subset of seven of the ten groups were also cultured in the presence of the reverse peptide, A $\beta$ 40-1. As a positive control, additional cells were cultured with the T cell mitogen, PHA, and, as a negative control, cells were cultured without added peptide.

Lymphocytes from a majority of the animals proliferated in response to PHA. There were no significant responses to the A $\beta$ 40-1 reverse peptide. Cells from animals immunized with the larger aggregated A $\beta$  peptides, AN1792, rodent A $\beta$ 1-42 and AN1528

when proliferated robustly ~~when~~ stimulated with A $\beta$ 1-40 with the highest cpm in the recipients of AN1792. One animal in each of the groups immunized with A $\beta$ 1-12 conjugate, A $\beta$ 13-28 conjugate and A $\beta$ 25-35 proliferated in response to A $\beta$ 1-40. The remaining groups receiving A $\beta$ 1-5 conjugate, A $\beta$ 33-42 conjugate pBx6 or PBS had no animals with an A $\beta$ -stimulated response. These results are summarized in Table 7 below.

Table 7			
Immunogen	Conjugate	A $\beta$ Amino Acids	Responders
A $\beta$ 1-5	Yes	5-mer	0/7
A $\beta$ 1-12	Yes	12-mer	1/8
A $\beta$ 13-28	Yes	16-mer	1/9
A $\beta$ 25-35		11-mer	1/9
A $\beta$ 33-42	Yes	10-mer	0/10
A $\beta$ 1-40		40-mer	5/8
A $\beta$ 1-42		42-mer	9/9
r A $\beta$ 1-42		42-mer	8/8
pBx6			0/8
PBS		0-mer	0/8

These results show that AN1792 and AN1528 stimulate strong T cell responses, most likely of the CD4+ phenotype. The absence of an A $\beta$ -specific T cell response in animals immunized with A $\beta$ 1-5 is not surprising since peptide epitopes recognized by CD4+ T cells are usually about 15 amino acids in length, although shorter peptides can sometimes function with less efficiency. Thus the majority of helper T cell epitopes for the four conjugate peptides are likely to reside in the IgG conjugate partner, not in the A $\beta$  region. This hypothesis is supported by the very low incidence of proliferative responses for animals in each of these treatment groups. Since the A $\beta$ 1-5 conjugate was effective at significantly reducing the level of A $\beta$  in the brain, in the apparent absence of A $\beta$ -specific T cells, the key effector immune response induced by immunization with this peptide appears to be antibody.

Lack of T-cell and low antibody response from fusion peptide pBx6, encompassing APP amino acids 592-695 including all of the A $\beta$  residues may be due to the poor immunogenicity of this particular preparation. The poor immunogenicity of the A $\beta$ 25-35 aggregate is likely due to the peptide being too small to be likely to contain a

to help

response.

good T cell epitope ~~to help~~ the induction of an antibody ~~response~~. It is anticipated that conjugation of this peptide to a carrier protein would render it more immunogenic.

#### V. Preparation of Polyclonal Antibodies for Passive Protection

- 5 125 non-transgenic mice were immunized with A $\beta$ , plus adjuvant, and euthanized at 4-5 months. Blood was collected from immunized mice. IgG was separated from other blood components. Antibody specific for the immunogen may be partially purified by affinity chromatography. An average of about 0.5-1 mg of immunogen-specific antibody is obtained per mouse, giving a total of 60-120 mg.

10

#### VI. Passive Immunization with Antibodies to A $\beta$

- Groups of 7-9 month old PDAPP mice each were injected with 0.5 mg in PBS of polyclonal anti-A $\beta$  or specific anti-A $\beta$  monoclonals as shown below. All antibody preparations were purified to have low endotoxin levels. Monoclonals can be prepared  
15 against a fragment by injecting the fragment or longer form of A $\beta$  into a mouse, preparing hybridomas and screening the hybridomas for an antibody that specifically binds to a desired fragment of A $\beta$  without binding to other nonoverlapping fragments of A $\beta$ .

Table 8

Antibody	Epitope
2H3	A $\beta$ 1-12
10D5	A $\beta$ 1-12
266	A $\beta$ 13-28
21F12	A $\beta$ 33-42
Mouse polyclonal anti-human A $\beta$ 42	Anti-Aggregated A $\beta$ 42

20

Mice were injected ip as needed over a 4 month period to maintain a circulating antibody concentration measured by ELISA titer of greater than 1/1000 defined by ELISA

## immunogen.

to A $\beta$ 42 or other ~~immunogen~~. Titers were monitored as above and mice were euthanized at the end of 6 months of injections. Histochemistry, A $\beta$  levels and toxicology were performed post mortem. Ten mice were used per group. Additional studies of passive immunization are described in Examples XI and XII below.

5

## VII. Comparison of Different Adjuvants

This example compares CFA, alum, an oil-in water emulsion and MPL for capacity to stimulate an immune response.

10

### A. Materials and Methods

#### 1. Study Design

One hundred female Hartley strain six-week old guinea pigs, obtained from Elm Hill Breeding Laboratories, Chelmsford, MA, were sorted into ten groups to be immunized with AN1792 or a palmitoylated derivative thereof combined with various adjuvants. Seven groups received injections of AN1792 (33  $\mu$ g unless otherwise specified) combined with a) PBS, b) Freund's adjuvant, c) MPL, d) squalene, e) MPL/squalene f) low dose alum, or g) high dose alum (300  $\mu$ g AN1792). Two groups received injections of a palmitoylated derivative of AN1792 (33  $\mu$ g) combined with a) PBS or b) squalene. A final, tenth group received PBS alone without antigen or additional adjuvant. For the group receiving Freund's adjuvant, the first dose was emulsified with CFA and the remaining four doses with IFA. Antigen was administered at a dose of 33  $\mu$ g for all groups except the high dose alum group, which received 300  $\mu$ g of AN1792. Injections were administered intraperitoneally for CFA/IFA and intramuscularly in the hind limb quadriceps alternately on the right and left side for all other groups. The first three doses were given on a biweekly schedule followed by two doses at a monthly interval). Blood was drawn six to seven days following each immunization, starting after the second dose, for measurement of antibody titers.

30

#### 2. Preparation of Immunogens

Two mg A $\beta$ 42 (California Peptide, Lot ME0339) was added to 0.9 ml of deionized water and the mixture was vortexed to generate a relatively uniform suspension.



10X

A 100  $\mu$ l aliquot of ~~10X~~ <sup>10X</sup> PBS (1X PBS, 0.15 M NaCl, 0.01 M sodium phosphate, pH 7.5) was added. The suspension was vortexed again and incubated overnight at 37°C for use the next day. Unused A $\beta$ 1-42 was stored with desiccant as a lyophilized powder at -20°C.

A palmitoylated derivative of AN1792 was prepared by coupling palmitic anhydride, dissolved in dimethyl formamide, to the amino terminal residue of AN1792 prior to removal of the nascent peptide from the resin by treatment with hydrofluoric acid.

To prepare immunogenic formulation doses with Complete Freund's adjuvant (CFA) (group 2), 33  $\mu$ g of AN1792 in 200  $\mu$ l PBS was emulsified 1:1 (vol:vol) with CFA in a final volume of 400  $\mu$ l for the first immunization. For subsequent immunizations, the antigen was similarly emulsified with Incomplete Freund's adjuvant (IFA).

To prepare formulation doses with MPL for groups 5 and 8, lyophilized powder (Ribi ImmunoChem Research, Inc., Hamilton, MT) was added to 0.2% aqueous triethylamine to a final concentration of 1 mg/ml and vortexed. The mixture was heated to 65 to 70°C for 30 sec to create a slightly opaque uniform suspension of micelles. The solution was freshly prepared for each set of injections. For each injection in group 5, 33  $\mu$ g of AN1792 in 16.5  $\mu$ l PBS, 50  $\mu$ g of MPL (50  $\mu$ l) and 162  $\mu$ l of PBS were mixed in a borosilicate tube immediately before use.

To prepare formulation doses with the low oil-in-water emulsion, AN1792 in PBS was added to 5% squalene, 0.5% Tween 80, 0.5% Span 85 in PBS to reach a final single dose concentration of 33  $\mu$ g AN1792 in 250  $\mu$ l (group 6). The mixture was emulsified by passing through a two-chambered hand-held device 15 to 20 times until the emulsion droplets appeared to be about equal in diameter to a 1.0  $\mu$ m diameter standard latex bead when viewed under a microscope. The resulting suspension was opalescent, milky white. The emulsions were freshly prepared for each series of injections. For group 8, MPL in 0.2% triethylamine was added at a concentration of 50  $\mu$ g per dose to the squalene and detergent mixture for emulsification as noted above. For the palmitoyl derivative (group 7), 33  $\mu$ g per dose of palmitoyl-NH-A $\beta$ 1-42 was added to squalene and vortexed. Tween 80 and Span 85 were then added with vortexing. This mixture was added to PBS to reach final concentrations of 5% squalene, 0.5% Tween 80, 0.5% Span 85 and the mixture was emulsified as noted above.

To prepare formulation doses with alum (groups 9 and 10), AN1792 in PBS was added to Alhydrogel (aluminum hydroxide gel, Accurate, Westbury, NY) to reach

gave modest titers ~~when~~ presented in an o/w emulsion with GMTs of 340 and 105 for the third and fourth bleeds. The highest antibody titers were generated with Freund's adjuvant with a peak GMT of about 87,000, a value almost 30-fold greater than the GMTs of the next two most potent formulations, MPL and high dose AN1792/alum.

- 5           The most promising adjuvants identified in this study are MPL and alum. Of these two, MPL appears preferable because a 10-fold lower antigen dose was required to generate the same antibody response as obtained with alum. The response can be increased by increasing the dose of antigen and /or adjuvant and by optimizing the immunization schedule. The o/w emulsion was a very weak adjuvant for AN1792 and
- 10       adding an o/w emulsion to MPL adjuvant diminished the intrinsic adjuvant activity of MPL alone.

## 2. A $\beta$ Levels in the Brain

- At about 14 weeks the guinea pigs were deeply anesthetized, the cerebrospinal
- 15   fluid (CSF) was drawn and brains were excised from animals in a subset of the groups, those immunized with Freund's adjuvant (group 2), MPL (group 5), alum with a high dose, 300  $\mu$ g, of AN1792 (group 10) and the PBS immunized control group (group 3). To measure the level of A $\beta$  peptide, one hemisphere was dissected and homogenates of the hippocampal, cortical, and cerebellar regions were prepared in 5 M guanidine. These were
- 20   diluted and quantitated by comparison to a series of dilutions of A $\beta$  standard protein of known concentrations in an ELISA format. The levels of A $\beta$  protein in the hippocampus, the cortex and the cerebellum were very similar for all four groups despite the wide range of antibody responses to A $\beta$  elicited by these formulations. Mean A $\beta$  levels of about 25 ng/g tissue were measured in the hippocampus, 21 ng/g in the cortex, and 12 ng/g in the
- 25   cerebellum. Thus, the presence of a high circulating antibody titer to A $\beta$  for almost three months in some of these animals did not alter the total A $\beta$  levels in their brains. The levels of A $\beta$  in the CSF were also quite similar between the groups. The lack of large effect of AN1792 immunization on endogenous A $\beta$  indicates that the immune response is focused on pathological formations of A $\beta$ .

30

## VIII. Immune Response to Different Adjuvants in Mice

female

for this

09/724570

Six-week old ~~male~~ Swiss Webster mice were used ~~for~~ <sup>for this</sup> study with 10-13 animals per group. Immunizations were given on days 0, 14, 28, 60, 90 and 20 administered subcutaneously in a dose volume of 200  $\mu$ l. PBS was used as the buffer for all formulations. Animals were bleed seven days following each immunization starting 5 after the second dose for analysis of antibody titers by ELISA. The treatment regime of each group is summarized in Table 9.

[REMAINDER OF PAGE INTENTIONALLY BLANK]

antibodies

shown

The ELISA titers of ~~antibodies~~ against A $\beta$ 42 in each group are ~~shown~~ in Table 10 below.

Table 10.

Geometric Mean Antibody Titers					
Week of Bleed					
Treatment Group	2.9	5.0	8.7	12.9	16.7
1	248	1797	2577	6180	4177
2	598	3114	3984	5287	6878
3	1372	5000	7159	12333	12781
4	1278	20791	14368	20097	25631
5	3288	26242	13229	9315	23742
6	61	2536	2301	1442	4504
7	37	395	484	972	2149
8	25	25	25	25	25
9	25	183	744	952	1823
10	25	89	311	513	817
11	29	708	2618	2165	3666
12	198	1458	1079	612	797
13	38	433	566	1080	626
14	104	541	3247	1609	838
15	212	2630	2472	1224	1496
16	183	2616	6680	2085	1631
17	28	201	375	222	1540
18	31699	15544	23095	6412	9059
19	63	243	554	299	441

The table shows that the highest titers were obtained for groups 4, 5 and 18, in which the adjuvants were 125  $\mu$ g MPL, 50  $\mu$ g MPL and QS-21 plus MPL.

#### 5 IX. Therapeutic Efficacy of Different Adjuvants

A therapeutic efficacy study was conducted in PDAPP transgenic mice with a set of adjuvants suitable for use in humans to determine their ability to potentiate immune responses to A $\beta$  and to induce the immune-mediated clearance of amyloid deposits in the brain.

- 10 One hundred eighty male and female, 7.5- to 8.5-month old heterozygous PDAPP transgenic mice were obtained from Charles River Laboratories. The mice were sorted into nine groups containing 15 to 23 animals per group to be immunized with AN1792 or

with

distributed

09/724570

AN1528 combined with various adjuvants. Animals were distributed to match the gender, age, and parentage of the animals within the groups as closely as possible. The adjuvants included alum, MPL, and QS-21, each combined with both antigens, and Freund's adjuvant (FA) combined with only AN1792. An additional group was immunized with AN1792 formulated in PBS buffer plus the preservative thimerosal without adjuvant. A ninth group was immunized with PBS alone as a negative control.

Preparation of aggregated A $\beta$  peptides: human A $\beta$ 1-40 (AN1528; California Peptides Inc., Napa, CA; Lot ME0541) and human A $\beta$ 1-42 (AN1792; California Peptides Inc., Lot ME0439) peptides were freshly solubilized for the preparation of each set of injections from lyophilized powders that had been stored desiccated at -20°C. For this purpose, two mg of peptide were added to 0.9 ml of deionized water and the mixture was vortexed to generate a relatively uniform solution or suspension. AN1528 was soluble at this step, in contrast to AN1792. A 100  $\mu$ l aliquot of 10X PBS (1X PBS: 0.15 M NaCl, 0.01 M sodium phosphate, pH 7.5) was then added at which point AN1528 began to precipitate. The suspensions were vortexed again and incubated overnight at 37°C for use the next day.

To prepare formulation doses with alum (Groups 1 and 5), A $\beta$  peptide in PBS was added to Alhydrogel (two percent aqueous aluminum hydroxide gel, Sargeant, Inc., Clifton, NJ) to reach concentrations of 100  $\mu$ g A $\beta$  peptide per 1 mg of alum. 10X PBS was added to a final dose volume of 200  $\mu$ l in 1X PBS. The suspension was then gently mixed for approximately 4 hr at RT prior to injection.

To prepare formulation doses for with MPL (Groups 2 and 6), lyophilized powder (Ribi ImmunoChem Research, Inc., Hamilton, MT; Lot 67039-E0896B) was added to 0.2% aqueous triethylamine to a final concentration of 1 mg/ml and vortexed. The mixture was heated to 65 to 70°C for 30 sec to create a slightly opaque uniform suspension of micelles. The solution was stored at 4°C. For each set of injections, 100  $\mu$ g of peptide per dose in 50  $\mu$ l PBS, 50  $\mu$ g of MPL per dose (50  $\mu$ l) and 100  $\mu$ l of PBS per dose were mixed in a borosilicate tube immediately before use.

To prepare formulation doses with QS-21 (Groups 3 and 7), lyophilized powder (Aquila, Framingham, MA; Lot A7018R) was added to PBS, pH 6.6-6.7 to a final concentration of 1 mg/ml and vortexed. The solution was stored at -20°C. For each set of injections, 100  $\mu$ g of peptide per dose in 50  $\mu$ l PBS, 25  $\mu$ g of QS-21 per dose in 25  $\mu$ l

PBS

tube

PBS and 125  $\mu$ l of ~~PBS~~ per dose were mixed in a borosilicate ~~tube~~ immediately before use.

To prepare formulation doses with Freund's Adjuvant (Group 4), 100  $\mu$ g of AN1792 in 200  $\mu$ l PBS was emulsified 1:1 (vol:vol) with Complete Freund's Adjuvant (CFA) in a final volume of 400  $\mu$ l for the first immunization. For subsequent immunizations, the antigen was similarly emulsified with Incomplete Freund's Adjuvant (IFA). For the formulations containing the adjuvants alum, MPL or QS-21, 100  $\mu$ g per dose of AN1792 or AN1528 was combined with alum (1 mg per dose) or MPL (50  $\mu$ g per dose) or QS-21 (25  $\mu$ g per dose) in a final volume of 200  $\mu$ l PBS and delivered by subcutaneous inoculation on the back between the shoulder blades. For the group receiving FA, 100  $\mu$ g of AN1792 was emulsified 1:1 (vol:vol) with Complete Freund's adjuvant (CFA) in a final volume of 400  $\mu$ l and delivered intraperitoneally for the first immunization, followed by a boost of the same amount of immunogen in Incomplete Freund's adjuvant (IFA) for the subsequent five doses. For the group receiving AN1792 without adjuvant, 10  $\mu$ g AN1792 was combined with 5  $\mu$ g thimerosal in a final volume of 50  $\mu$ l PBS and delivered subcutaneously. The ninth, control group received only 200  $\mu$ l PBS delivered subcutaneously. Immunizations were given on a biweekly schedule for the first three doses, then on a monthly schedule thereafter on days 0, 16, 28, 56, 85 and 112. Animals were bled six to seven days following each immunization starting after the second dose for the measurement of antibody titers. Animals were euthanized approximately one week after the final dose. Outcomes were measured by ELISA assay of A $\beta$  and APP levels in brain and by immunohistochemical evaluation of the presence of amyloid plaques in brain sections. In addition, A $\beta$ -specific antibody titers, and A $\beta$ -dependent proliferative and cytokine responses were determined.

Table 10 shows that the highest antibody titers to A $\beta$ 1-42 were elicited with FA and AN1792, titers which peaked following the fourth immunization (peak GMT: 75,386) and then declined by 59% after the final, sixth immunization. The peak mean titer elicited by MPL with AN1792 was 62% lower than that generated with FA (peak GMT: 28,867) and was also reached early in the immunization scheme, after 3 doses, followed by a decline to 28% of the peak value after the sixth immunization. The peak mean titer generated with QS-21 combined with AN1792 (GMT: 1,511) was about 5-fold lower than obtained with MPL. In addition, the kinetics of the response were slower, since an

- immunization response.
- additional immunization was required to reach the peak response. Titers generated by alum-bound AN1792 were marginally greater than those obtained with QS-21 and the response kinetics were more rapid. For AN1792 delivered in PBS with thimerosal the frequency and size of titers were barely greater than that for PBS alone. The peak titers generated with MPL and AN1528 (peak GMT 3099) were about 9-fold lower than those with AN1792. Alum-bound AN1528 was very poorly immunogenic with low titers generated in only some of the animals. No antibody responses were observed in the control animals immunized with PBS alone.

Table 11

Geometric Mean Antibody Titers <sup>a</sup>					
Week of Bleed					
Treatment	3.3	5.0	9.0	13.0	17.0
Alum/	102	1,081	2,366	1,083	572
AN1792	(12/21) <sup>b</sup>	(17/20)	(21/21)	(19/21)	(18/21)
MPL/	6241	28,867	1,1242	5,665	8,204
AN1792	(21/21)	(21/21)	(21/21)	(20/20)	(20/20)
QS-21/	30	227	327	1,511	1,188
AN1792	(1/20)	(10/19)	(10/19)	(17/18)	(14/18)
CFA/	10,076	61,279	75,386	41,628	30,574
AN1792	(15/15)	(15/15)	(15/15)	(15/15)	(15/15)
Alum/	25	33	39	37	31
AN1528	(0/21)	(1/21)	(3/20)	(1/20)	(2/20)
MPL/	184	2,591	1,653	1,156	3,099
AN1528	(15/21)	(20/21)	(21/21)	(20/20)	(20/20)
QS-21/	29	221	51	820	2,994
AN1528	(1/22)	(13/22)	(4/22)	(20/22)	(21/22)
PBS plus	25	33	39	37	47
Thimerosal	(0/16)	(2/16)	(4/16)	(3/16)	(4/16)
PBS	25	25	25	25	25
	(0/16)	(0/16)	(0/15)	(0/12)	(0/16)

## 10 Footnotes:

<sup>a</sup> Geometric mean antibody titers measured against A $\beta$ 1-42<sup>b</sup> Number of responders per group

Example

animals,

and Freund's adjuvant in ~~Example~~ VI relative to PBS treated ~~animals~~ these type of effects are expected from Freund's adjuvant treatment and the accompanying peritonitis and do not indicate any adverse effects from AN1792 treatment. Although not part of the toxicological evaluation, PDAPP mouse brain pathology was extensively examined as part of the efficacy endpoints. No sign of treatment related adverse effect on brain morphology was noted in any of the studies. These results indicate that AN1792 treatment is well tolerated and at least substantially free of side effects.

#### XI. Therapeutic Treatment with Anti-A $\beta$ antibodies

The experiments described in this section were carried out in order to test the abilities of various monoclonal and polyclonal antibodies against A $\beta$  to inhibit accumulation of A $\beta$  in the brain of heterozygotic transgenic mice.

##### A. Study 1

##### 1. Study Design

Sixty male and female, heterozygous PDAPP transgenic mice, 8.5 to 10.5 months of age were obtained from Charles River Laboratory. The mice were sorted into six groups to be treated with various antibodies directed to A $\beta$ . Animals were distributed to match the gender, age, parentage and source of the animals within the groups as closely as possible. As shown in Table 12, the antibodies included four murine A $\beta$ -specific monoclonal antibodies, 2H3 (directed to A $\beta$  residues 1-12), 10D5 (directed to A $\beta$  residues 1-16), 266 (directed to A $\beta$  residues 13-28 and binds to monomeric but not to aggregated AN1792), 21F12 (directed to A $\beta$  residues 33-42). A fifth group was treated with an A $\beta$ -specific polyclonal antibody fraction (raised by immunization with aggregated AN1792). The negative control group received the diluent, PBS, alone without antibody.

The monoclonal antibodies were injected at a dose of about 10 mg/kg (assuming that the mice weighed 50 g). Injections were administered intraperitoneally every seven days on average to maintain anti-A $\beta$  titers above 1000. Although lower titers were measured for mAb 266 since it does not bind well to the aggregated AN1792 used as the capture antigen in the assay, the same dosing schedule was maintained for this group. The group receiving monoclonal antibody 2H3 was discontinued within the first three weeks since the antibody was cleared too rapidly in vivo. Animals were bled prior to each dosing



for the measurement of antibody titers. Treatment was continued over a six-month period for a total of 196 days. Animals were euthanized one week after the final dose.

Table 12

<u>EXPERIMENTAL DESIGN OF STUDY 006</u>				
Treatment Group	N <sup>a</sup>	Treatment Antibody	Antibody Specificity	Antibody Isotype
1	9	none (PBS alone)	NA <sup>b</sup>	NA
2	10	Polyclonal	A $\beta$ 1-42	mixed
3	0	mAb <sup>c</sup> 2H3	A $\beta$ 1-12	IgG1
4	8	mAb 10D5	A $\beta$ 1-16	IgG1
5	6	mAb 266	A $\beta$ 13-28	IgG1
6	8	mAb 21F12	A $\beta$ 33-42	IgG2a

## Footnotes

- a. Number of mice in group at termination of the experiment. All groups started with 10 animals per group.  
 b. NA: not applicable  
 c. mAb: monoclonal antibody

## 2. Materials and Methods

## a. Preparation of the Antibodies

The anti-A $\beta$  polyclonal antibody was prepared from blood collected from two groups of animals. The first group consisted of 100 female Swiss Webster mice, 6 to 8 weeks of age. They were immunized on days 0, 15, and 29 with 100  $\mu$ g of AN1792 combined with CFA/IFA. A fourth injection was given on day 36 with one-half the dose of AN1792. Animals were exsanguinated upon sacrifice at day 42, serum was prepared and the sera were pooled to create a total of 64 ml. The second group consisted of 24 female mice isogenic with the PDAPP mice but nontransgenic for the human APP gene, 6 to 9 weeks of age. They were immunized on days 0, 14, 28 and 56 with 100  $\mu$ g of AN1792 combined with CFA/IFA. These animals were also exsanguinated upon sacrifice at day 63, serum was prepared and pooled for a total of 14 ml. The two lots of sera were pooled. The antibody fraction was purified using two sequential rounds of precipitation

ammonium

was

with 50% saturated ammonium sulfate. The final precipitate was dialyzed against PBS and tested for endotoxin. The level of endotoxin was less than 1 EU/mg.

The anti-A $\beta$  monoclonal antibodies were prepared from ascites fluid. The fluid was first delipidated by the addition of concentrated sodium dextran sulfate to ice-cold ascites fluid by stirring on ice to reach a final concentration of 0.238%. Concentrated CaCl<sub>2</sub> was then added with stirring to reach a final concentration of 64mM. This solution was centrifuged at 10,000 x g and the pellet was discarded. The supernatant was stirred on ice with an equal volume of saturated ammonium sulfate added dropwise. The solution was centrifuged again at 10,000 x g and the supernatant was discarded. The pellet was resuspended and dialyzed against 20 mM Tris-HCl, 0.4 M NaCl, pH 7.5. This fraction was applied to a Pharmacia FPLC Sepharose Q Column and eluted with a reverse gradient from 0.4 M to 0.275 M NaCl in 20 mM Tris-HCl, pH 7.5.

The antibody peak was identified by absorbance at 280 nm and appropriate fractions were pooled. The purified antibody preparation was characterized by measuring the protein concentration using the BCA method and the purity using SDS-PAGE. The pool was also tested for endotoxin. The level of endotoxin was less than 1 EU/mg, titers, titers less than 100 were arbitrarily assigned a titer value of 25.

### 3. A $\beta$ and APP Levels in the Brain:

Following about six months of treatment with the various anti-A $\beta$  antibody preparations, brains were removed from the animals following saline perfusion. One hemisphere was prepared for immunohistochemical analysis and the second was used for the quantitation of A $\beta$  and APP levels. To measure the concentrations of various forms of beta amyloid peptide and amyloid precursor protein (APP), the hemisphere was dissected and homogenates of the hippocampal, cortical, and cerebellar regions were prepared in 5M guanidine. These were serially diluted and the level of amyloid peptide or APP was quantitated by comparison to a series of dilutions of standards of A $\beta$  peptide or APP of known concentrations in an ELISA format.

The levels of total A $\beta$  and of A $\beta$ 1-42 measured by ELISA in homogenates of the cortex, and the hippocampus and the level of total A $\beta$  in the cerebellum are shown in Tables 11, 12, and 13, respectively. The median concentration of total A $\beta$  for the control group, inoculated with PBS, was 3.6-fold higher in the hippocampus than in the cortex

ng/g hippocampal

17,818 ng/g

(median of 63,389 ng/g hippocampal tissue compared to 17,818 ng/g for the cortex). The median level in the cerebellum of the control group (30.6 ng/g tissue) was more than 2,000-fold lower than in the hippocampus. These levels are similar to those that we have previously reported for heterozygous PDAPP transgenic mice of this age (Johnson-Woods et al., 1997).

For the cortex, one treatment group had a median A $\beta$  level, measured as A $\beta$ 1-42, which differed significantly from that of the control group ( $p < 0.05$ ), those animals receiving the polyclonal anti-A $\beta$  antibody as shown in Table 13. The median level of A $\beta$ 1-42 was reduced by 65%, compared to the control for this treatment group. The median levels of A $\beta$ 1-42 were also significantly reduced by 55% compared to the control in one additional treatment group, those animals dosed with the mAb 10D5 ( $p = 0.0433$ ).

[REMAINDER OF THIS PAGE INTENTIONALLY BLANK]